

INDEX

- Abscesses, 243–44, 244f
Acetabular fractures, diagnosis of, 392, 393f
Acetabulum, abnormalities, 384f
Acetylsalicylic acid (aspirin), 414
ACS. *See* Autologous conditioned serum
American Association of Equine Practitioners, 81t
Anatomy, functional, of musculoskeletal system
 antebrachium, 22–23
 arm, shoulder, 25, 27f–28f
 carpus, 18, 20, 20f
 crus, 43, 44f–45f
 cubital joint, 23, 23f–24f, 25
 digit, fetlock, 3, 4f
 dorsal aspect, 18
 extensor muscles, 20f, 23, 24f
 fetlock, 14, 15f–17f
 of foot, 3
 hindlimb, 32, 35f, 36
 hip, 51–56, 52f–55f
 hoof wall, sole, frog, 4–9, 6f–9f
 lateral aspect, 20, 20f–21f
 medial aspect, 21, 21f
 metacarpus, 14, 18
 nomenclature, usage, 3
 palmar aspect, 18, 19f, 21–22, 22f
 pastern, 11f–12f, 13–14
 positional, directional terms for, 4f
 stay apparatus, 31–32
 stifle, 43, 44f–49f, 46
 tarsal joint, 40–43, 41f–42f
 thigh, 51–56, 52f–55f
Anesthesia. *See also* Joint anesthesia;
 Perineural anesthesia
 bicipital bursa, 144–45, 145f
 calcaneal bursa, 144, 144f
 carpal sheath, 142–43
 cunean bursa, 145–46
 DFTS, 141–42, 141f–142f
 DIP, 126–27, 127f–128f
 DIT, 136–37, 137f
 in ECR, 143
 FP, 139, 193f
 hip joint, 140, 140f
 intrasynovial, 124–26, 125t
 LFT, 139–40, 140f
 MFT, 138–39, 138f
 middle carpal joint, 131
 for navicular bone, 143–44, 143f–144f
 PIP, 127–28, 129f
 radiocarpal joint, 131–32
 shoulder joint, 134–36, 135f
 SI, 140–41, 141f
 tarsal sheath, 143
 TC, 137–38
 tendon sheath, 141–43, 141f–142f
 TMT, 136, 136f
 trochanteric bursa, 145, 146f
Antebrachium
 anatomy, nomenclature of, 22–23
 medial cutaneous blocks of, 121, 121f
Antimicrobials, for synovial wound
 treatment of, 437t, 439t
Arm. *See also* Forearm
 anatomy, nomenclature of, 25, 27f–28f
 medial view of, 26f
Arquel. *See* Meclofenamic acid
Arterial supply, to forelimb digit, 15f
Arthritis. *See also* Osteoarthritis
 infectious, of hip joint, 383–84, 384f
 septic, 164f, 166f
Arthrodesis, of fetlock joint, 297f
Artifacts, for MRI, 209–12, 211f
Aspirin. *See* Acetylsalicylic acid
ASU. *See* Avocado and soybean unsaponified extracts
Asymmetrical displacement, 258f, 261f
Atlas, 58f
Autologous conditioned serum (ACS),
 intrasynovial, 411
Avocado and soybean unsaponified
 extracts (ASU), 417
Avulsion fracture, 228f
Axial components
 caudal vertebrae, 60
 cervical vertebrae, 58–59, 58f
 ligamentum nuchae, 61
 lumbar vertebrae, 58f, 59
 sacroiliac region, 60–61, 61f
 thoracic vertebrae, 58f, 59
 vertebral articulations of, 60
 vertebral column, 56, 58f
Axis, 58f

- Back, palpation, manipulation of, for lameness assessment, 104, 107f–108f
- Banamine®. *See* Flunixin Meglumine
- Bandages and Bandaging, types of, 407
- Basisesamoid block (High PD), 116–17, 116f
- Bicipital bursa anesthesia, 144–45, 145f
- Bicipital bursitis (intertubercular bursa)
diagnosis of, 332–33, 333f
treatment of, 333, 334f
- Bisphosphonates, 406
- Bizolin®. *See* Phenylbutazone
- Bone marrow mesenchymal stem cell therapy, 411
- Bone response patterns, 161–63, 163f
- Bones. *See also specific bones*
cortical changes in, 160, 161f
palpation of, 93f
of pelvic limb, 34f
radiographic interpretation of, 159–63, 160f–163f
radiology of, 159–63, 160f–163f
radiopharmaceutical uptake of, 192
of thoracic limb, 5f
ultrasound evaluation of, 180, 182, 191f
- Brass band, 257f
- Breakover-heel lift, 71f
- Breakover-toe pivot, 71f
- Bruising, soft tissue, 430f
- Bucked shins, 299–301, 300f–301f
- Bursa anesthesia
bicipital, 144–45, 145f
calcaneal, 144, 144f
cunean, 145–46
navicular, 143–44, 143f–144f
trochanteric, 145, 146f
- Bursitis, of elbow
clinical signs of, 329, 329f
treatment of, 330
- Butatron™. *See* Phenylbutazone
- Butazolidin®. *See* Phenylbutazone
- Butequine®. *See* Phenylbutazone
- Buttress foot, 235f
- Calcaneal bursa
anesthesia, 144, 144f
infection in, puncture and, 430f
- Calcaneal bursitis, 352–53
- Calcaneus, luxation of, of SDFT, 350–52, 351f–353f
- Cancellous bone, signal intensity of, 208t
- Canker
clinical signs of, 244–45, 244f
treatment of, 245f
- Cannon bone
fractures of, 301–4, 302f, 304f–305f
imaging of, 173, 174f
- Canter, example of, 69f
- Capital physal fractures, of femoral head, 379–80, 379f
- Capped hock, 352–53
- Capsulitis, MFT, 375
- Carpal bones
fractures of, 323f
osteochondral fractures of, 319f
- Carpal canal swelling, 322f
- Carpal flexion tests, manipulation of, 110
- Carpal ligaments, 20f
medial view of, 21f
- Carpal sheath
anesthesia, 142–43
tenosynovitis, 321–23, 322f–323f
- Carprofen (Zenecarp, Rimadyl®), 415
- Carpus
anatomy, nomenclature of, 18, 20, 20f
deep dissection of, 20f
dorsal view of, 19f
lateral view of, 21f
minimum image acquisition counts in, 193t
MRI protocol for, 210t
OA of
clinical signs of, 320, 320f
diagnosis of, 320, 321f
treatment of, 320–21
osteochondroma of, 322f
palmaromedial view of, with flexor retinaculum cut, 22f
palpation, manipulation of, for lameness assessment, 93, 95f–96f
- Cartilage
collateral, ossification of, 241–43, 242f
signal intensity of, 208t
- Caudal cruciate test, 103f
- Caudal splint, 435f
- Caudal vertebrae, 60
- CDE. *See* Common digital extensor tendon rupture
- Center of pressure, for shoes, 420–21, 420f–422f
- Central tarsal bones, SLAB, sagittal fractures of, 347–48, 348f
- Cervical facet joint osteoarthritis, treatment of, 403
- Cervical vertebrae, anatomy of, 58–59, 58f
- CH. *See* Collagen hydrolysate
- Chip fractures. *See* Osteochondral fractures
- Chondroitin sulfate (CS), 416
with GLN, 416
- Club foot
clinical signs of, 253, 254f
diagnosis of, 253–54
treatment of, 254, 254f
- Coccygeal vertebrae, fractures of, 392–93, 393f
- Coffin bone, of fracture, 235–36
- Coffin joint, dorsolateral approach to, 127f
- Cold therapy, 407
- Collagen hydrolysate (CH), 417
- Collateral cartilage, ossification of, of distal phalanx, 241–43, 242f
- Collateral cruciate ligament injury
diagnosis of, 374, 374f
treatment of, 374
- Collateral desmitis, MRI diagnosis of, 214t
- Collateral ligaments
injuries to
diagnosis of, 233, 233f
treatment of, 233
palpation of, 89f
- Collateral sesamoidean ligament, approach, to fetlock, 131f
- Collateral sesamoidean ligaments (CSL), attachments of, 13f
- Comminuted fractures
CT of, 220f–221f
of femur, 378f
of P1, treatment of, 282, 282f
P2 fractures, 277, 278f
- Common digital extensor tendon rupture (CDE)
clinical signs of, 315, 316f
treatment of, 316
- Complete fractures
of MCIII, 303–4, 304f–305f
of MTIII, 303–4, 304f–305f
- Computed radiography
digital image display, 154, 154f
digital image storage, 154–55
in digital radiography systems, 151
direct digital radiography, 153–54, 153f
reading process for, 151, 152f, 153
- Computed tomography (CT)
of comminuted fractures, 220f–221f
of distal phalanx fractures, 220f
of fracture, 218f
imaging benefits of, 216–17
interpretation of, 217–19, 218f
for lameness assessment, indications, 219, 219f–221f
- Condylar fractures
of MCIII/MTIII
diagnosis of, 302–3, 302f
treatment of, 303
types of, 301f
- Corium
histological relationship to, 9f
relationship, to hoof, 7f
- Corns, 244f
treatment of, 243–44
- Coronary region, of hoof wall, three-dimensional dissection of, 8f
- Coronet band, dissection of, 8f
- Corrective trimming
distribution of force in, 421–22
for ease of movement, 421, 422f
foot motion and, during flight, 422–23

- heels in, 420f
 limb examination for, 418–19, 418f
 point of break-over in, 422f
 principles of, 419, 419t
 for protection, 423
 for support, 423
- Cortical bone
 changes in, 160, 161f
 signal intensity of, 208t
- Corticosteroids (CS)
 HA combinations with, 409–10
 intra-articular, 409t
 intralesional, 412–13
 intrasynovial therapy with, 408–9
- Counterirritation, from topical, local medications, 408
- Coxofemoral joint. *See* Hip joint
- Coxofemoral luxation
 clinical signs of, 380, 381f
 treatment of, 380–82
- Cranial cruciate test, 104f
- Crest fractures
 diagnosis of, 361–62
 treatment of, 361–62, 361f
- Cross sectional area (CSA), measurements, 175, 176f
- Cruciate ligament injury
 diagnosis of, 374, 374f
 treatment of, 374
- Crus (leg, gaskin)
 anatomy, nomenclature of, 43, 44f–45f
 caudal view of, 48f
 dorsal view of, 44f
 medial dissection of, 38f
 superficial dissection of, 45f
- CS. *See* Chondroitin sulfate; Corticosteroids
- CSA. *See* Cross sectional area
- CSL. *See* Collateral sesamoidean ligaments
- CT. *See* Computed tomography
- Cubital joint (elbow)
 anatomy, nomenclature of, 23, 23f–24f, 25
 bursitis of
 clinical signs of, 329, 329f
 treatment of, 330
 caudomedial view of, 24f
 cystic lesions of
 clinical signs of, 328–29
 diagnosis of, 328f, 329
 treatment of, 329
 immobilization of, 434, 435f
 joint anesthesia, 132–34, 135f
 lateral approaches to, 135f
 lateral view of, 27f
 minimum image acquisition counts in, 193t
 palpation of, 97f
- Cunean bursa anesthesia, 145–46
- Cystic lesions, of elbow
 diagnosis of, 328f, 329
 treatment of, 329
- DBLPN. *See* Deep branch of lateral plantar nerve block
- DDFT. *See* Deep digital flexor tendon
- Deep branch of lateral plantar nerve block (DBLPN), 123, 123f
- Deep digital flexor tendon (DDFT)
 attachments of, 13f
 drawing of, 12f
 imaging of, 172f, 174f, 176f–180f
 regions for, 182f
 injuries to, 232f
 treatment of, 232
 lacerations of, 441f
 palmaromedial view of, with flexor retinaculum cut, 22f
 pastern injuries
 treatment of, 285–86
 ultrasound of, 286f
 tendonitis, MRI diagnosis of, 214t
 tenotomy of, 262f
 transducers in, 183f–187f
 ultrasound assessment of, 172f, 174f, 176f–180f
- Degenerative suspensory ligament desmitis (DSL/D)
 clinical signs of, 311, 312f
 diagnosis of, 311–12
 treatment of, 312
- Delayed phase
 for enostosis-like lesion diagnosis of, 201f
 of nuclear scintigraphy, 197
 for palmar process fracture diagnosis of, 200f
 of rhabdomyolysis, 202f
 as scintigraphic sign, 197, 198f–202f, 199
 for stress fracture diagnosis of, 199f
 for tarsal bone fracture diagnosis of, 200f
- Dermal laminae, photomicrograph of, 9f
- Desmitis
 collateral, MRI diagnosis of, 214t
 DSL/D, 311–12, 312f
 of DSL/D
 clinical signs of, 283
 diagnosis of, 283–84, 284f
 MRI diagnosis of, 214t
 treatment of, 284
 intersesamoidean desmitis, MRI diagnosis of, 214t
 palmar annular, MRI diagnosis of, 214t
 proximal suspensory, 217f
 suspensory ligament, 310f
 degenerative, 311–14, 312f–314f
 diagnosis of, 310, 311f
 treatment of, 310
- DFTS. *See* Digital flexor tendon sheath
- Diaphyseal femoral fractures, clinical signs of, 377–78, 378f
- Diaphyseal tibial fractures
 clinical signs of, 359–60
- diagnosis of, 359f, 360
 treatment of, 360, 360f
- DICOM® software, 154
- Digital extensor tendon. *See* Common digital extensor tendon rupture
- Digital flexor synovial sheath, palpation of, 90f
- Digital flexor tendon sheath (DFTS)
 anesthesia, 141–42, 141f–142f
 distal approach to, 142f
 injection site, 142f
 proximal approach to, 141f
 swelling of, 298f
 tenosynovitis of
 clinical signs of, 298, 298f
 diagnosis of, 298–99, 298f
 treatment of, 299, 299f
- Digital image display, 154, 154f
- Digital image storage, 154–55
- Digital radiography systems
 computed radiography, 151
 reading process for, 151, 152f, 153
- Digits
 anatomy, 3, 4f
 forelimb, arterial supply to, 15f
 hindlimb, anatomy, nomenclature of, 32, 35f, 36
 sagittal section of, 11f
- Dimethyl sulfoxide (DMSO), 407–8
- DIP. *See* Distal interphalangeal joint
- Direct digital radiography, 153–54, 153f
 direct conversion, 153, 153f
 indirect conversion, 153–54, 153f
- Direct flat panel detector, 155f
- Direct/local pressure plus movement test, 113–14
- Discoepiphysitis, treatment of, 398
- Distal interphalangeal joint (DIP)
 anesthesia, 126–27, 127f–128f
 correct movement of, 420f
 OA of
 clinical signs of, 234, 234f
 treatment of, 235
- Distal intertarsal joint (DIT)
 anesthesia, 136–37, 137f
 medial, dorsolateral approaches to, 137f
- Distal limb flexion test, 91f, 109–10, 109f–110f
- Distal phalanx (P3, sidebone)
 attachments of, 13f
 displacement of, 258f
 fracture of, 236f–237f
 CT of, 220f
 diagnosis of, 236, 237f
 full bar shoes for, 238f
 treatment of, 237, 238t, 239
 types of, 238t
- ligaments, 10f
 ossification of, of collateral cartilage, 241–43, 242f
 rotation of, 260f
 subchondral cystic lesions of, clinical signs of, 240–41, 241f

- Distal sesamoid bone. *See* Navicular bone
- Distal sesamoid ligaments (DSLS), desmitis of
 clinical signs of, 283
 diagnosis of, 283–84, 284f
 MRI diagnosis of, 214t
 treatment of, 284
- Distal tarsal osteoarthritis
 clinical signs of, 344
 diagnosis of, 344, 344f–345f
 treatment of, 345
- Distribution of force, by shoes, trimming, 421–22
- DIT. *See* Distal intertarsal joint
- Dorsal aspect
 anatomy, nomenclature of, 18
 of tarsus, 36, 37f
- Dorsal spinous process, overriding and impingement of, generalizations of, 395
- Dorsoscapular ligaments
 attachments of, 26f, 29f, 31
 right, 28f
- Draft horses, occupation-related lameness in, 77t
- Draining tract, at shoulder joint, 193f
- Dressage horses, occupation-related lameness in, 77t
- Dropped elbow, in foals, 78f
- DSL. *See* Degenerative suspensory ligament desmitis
- DSLS. *See* Distal sesamoid ligaments
- Ease of movement, corrective shoeing, trimming for, 421, 422f
- Echo time (TE), 207
- ECR. *See* Extensor carpi radialis
- Elbow. *See* Cubital joint
- Elbow flexion tests, manipulation of, 110
- Eminence fractures, 277, 277f
- Endurance horses, occupation-related lameness in, 77t
- Enostosis-like lesion diagnosis of, delayed phase for, 201f
- Entheses, in soft tissue, 159, 159f
- Entosis-like lesion, MRI diagnosis of, 214t
- Epidermal laminae, photomicrograph of, 9f
- Equioxx®. *See* Firocoxib
- Equiphen®. *See* Phenylbutazone
- Equiproxen®. *See* Naproxen
- Equizone 100™. *See* Phenylbutazone
- Eventing horses, occupation-related lameness in, 77t
- Exercise, visual examination at, 107–8
- Extensor carpi radialis (ECR)
 damage, treatment of, 316–17, 317f
 palpation of, 95f
 tendon sheath anesthesia in, 143
- Extensor carpi ulnaris (Ulnaris lateralis), 21f
- Extensor muscles, anatomy, nomenclature of, 20f, 23, 24f
- Extensor process fracture, 238f
- Extracorporeal shock wave therapy, 408
- Facet joint osteoarthritis
 cervical, 402–3
 diagnosis of, 400, 401f–402f
 treatment of, 401
- Fast low angle shot (FLASH), 213f
- Fat-derived mesenchymal stem cell therapy, intrasynovial, 411
- Fats, signal intensity of, 208t
- Fatty acids, 417
- Femoral fractures, 377–78, 378f
 type II distal, 379f
- Femoral head
 abnormalities, 384f
 capital physal fractures of, 379–80, 379f
- Femoropatellar joint (FP)
 anatomy, 47–49
 anesthesia, 139, 193f
 cranial, lateral approaches to, 139f
- Femoropatellar osteochondrosis
 clinical signs of, 363–64
 diagnosis of, 364, 365f
 treatment of, 364
- Femorotibial joint
 anatomy of, 49–51
 opened capsule of, 49f
- Femur
 comminuted fractures of, 378f–379f
 palpation, manipulation of, for lameness assessment, 103–4
- Fetlock
 anatomy, 3, 4f
 anatomy, nomenclature of, 14, 15f–17f
 collateral sesamoidean ligament approach to, 131f
 distal palmar, plantar approach to, 132f
 dorsal injection site for, 133f
 hindlimb, anatomy, nomenclature of, 32, 35f, 36
 lameness assessment of, palpation, manipulation for, 89–90, 89f–90f
 MRI protocol for, 209t
 proximal palmar, plantar approach to, 130f
 sagittal section of, 11f
- Fetlock flexion test, 91f, 109–10, 109f–110f
- Fetlock joint, arthrodesis of, 297f
- Fetlock subchondral cystic lesions, treatment of, 294–95
- Fibrotic myopathy
 clinical signs of, 376, 376f
 of hindlimb, 101f, 376f
- myotomy for, 377f
 treatment of, 376–77
- Fibularis, lateral view of, 35f
- Film processing, for imaging, 151
- Firocoxib (Equioxx®), 415
- First aid kit, for fractures, 433t
- First aid treatments, 407
- FLASH. *See* Fast low angle shot
- Flexion tests, manipulation
 carpal, 110
 distal limb, 91f, 109–10, 109f–110f
 elbow, 110
 fetlock, 91f, 109–10, 109f–110f
 full limb forelimb, 112–13, 113f
 hindlimb, 112–13, 113f
 shoulder, upper forelimb, 110–11, 111f
 of stifle, 112, 113f
 tarsal, hock, 111, 112f
 types of, 108–9
- Flexor tendons. *See also* Deep digital flexor tendon; Digital flexor tendon sheath
 laceration of, 441f
 palpation, manipulation of, for lameness assessment, 92, 95f
- Fluids, signal intensity of, 208t
- Flunixin Meglumine (Banamine®), 414
- Foals
 dropped elbow in, 78f
 gastrocnemius muscle rupture in, 79f
 lameness evaluation of, 76–78, 78f
- Foot
 anatomy, 3
 buttress, 235f
 club, 253–55, 253f–254f
 flight, 73f
 imbalances
 clinical signs of, 251–52
 diagnosis of, 252, 252f
 treatment of, 252, 253f
 internal structures of, 9, 10f–11f
 lameness assessment in
 MRI for, 213t, 214f
 palpation, manipulation for, 85–87, 85f–87f
 ligaments of, 10f
 minimum image acquisition counts in, 193t
 motion of, during flight, 422–23
 MRI protocol for, 209t
 nerves of, 16f–17f
 penetrating injuries of
 diagnosis of, 248–49
 treatment of, 249
- Forearm
 caudomedial view of, 24f
 lameness assessment of, palpation, manipulation for, 96–97, 97f
 lateral view of, 21f
 proximal, medial view of, 26f
- Foreign bodies, ultrasound assessment of, 182, 185, 193f

- Forelimb
arterial supply to, 15f
full, flexion tests, manipulation of, 112–13, 113f
hindlimb lameness *vs.*, 74–75
lameness assessment in, 84f
lameness diagnosis of in, 72–73, 73b, 73f
palpation, manipulation of, for lameness assessment, 96–97, 97f
perineural anesthesia, 116–24, 116f–124f
upper, flexion tests, manipulation of, 110–11, 111f
- FP. *See* Femoropatellar joint
- Fractures. *See also specific types of fracture*
of cannon bone, 301–4, 302f, 304f–305f
of carpal bones, 323f
of coccygeal vertebrae, 392–93, 393f
of coffin bone, 235–36
CT of, 218f
of dorsal MC, 299–301, 300f–301f
of femur, 377–78, 378f–379f
first aid kit for, 433t
of humerus, diagnosis of, 330–31, 331f
of long bone, 432–36, 433t, 434f–436f
of MCIII, 301–3, 302f
of MCIII/MTIII, 303–4, 304f–305f
of navicular disease and syndromes, 230, 231f
diagnosis of, 230, 230f
treatment of, 230–31
of P1, 279f–280f, 281–83, 282f
of P2, 275–77, 276f–278f
of P3
clinical signs of, 236–37, 237f–238f, 238t
CT of, 220f
of patella, 364–67, 366f
of proximal sesamoid bones, treatment of, 288f, 290f–291f, 291
of proximal tibial physis, 362–63, 362f–363f
of radius
clinical signs of, 323–24
diagnosis of, 323–24, 324f
treatment of, 324, 325f
of sacrum, 392–93, 393f
of small metacarpus, 306–8, 307f–308f
of small metatarsus, 306–8, 307f–308f
of spinous process, 397
of supraglenoid tubercle, 337–38, 337f
of sustentaculum tali, 355f
of tarsal bone, 200f, 347–48, 348f
of tibial malleoli, 348–49
- of ulna
clinical signs of, 327, 327f
diagnosis of, 327, 327f
treatment of, 327, 327f–328f
- Frog
anatomy of, 4–9, 6f–9f
nail in, 86f
- Full bar shoes, for distal phalanx fracture, 238f
- Gaits, for lameness diagnosis of, fundamentals of, 67–69, 67f–69f
- Gallop, image of, 70f
- Gamma camera
positioning of, 196f
use of, 193
working of, 195
- Gas, in soft tissue, 158–59, 159f
- Gaskin. *See* Crus
- Gastrocnemius muscle, rupture of, in foals, 79f
- GE. *See* Gradient echo
- Genu. *See* Stifle
- Glucosamine (GLN), 416
chondroitin sulfate with, 416
- Gluteal atrophy, 106f
- Gradient echo (GE), 207
- Ground reaction force, 261f
- HA. *See* Hyaluronan
- Hallmarq Equine Limbscanner®, 205f
- Heel bulb, 88f
- Heel cracks
clinical signs of, 255–56
treatment of, 256, 256f–257f
- Heels
breakover-heel lift, 71f
in corrective trimming, 420f
sheared, 252f–253f
wedge of, 420f
- High palmar/high four-point block, 118, 119f
- High PD. *See* Basisesamoid block
- High plantar/sub-tarsal blocks, 122–23, 122f
- High two-point block, 120, 120f
- Hindlimb
anatomy, nomenclature of, 32, 35f, 36
digit, anatomy, nomenclature of, 32, 35f, 36
fetlock, anatomy, nomenclature of, 32, 35f, 36
fibrotic myopathy of, 101f, 376f
flexion tests, manipulation of, 112–13, 113f
forelimb lameness *vs.*, 74–75
lameness assessment in, 84f
lameness diagnosis of in, 73–74, 73t, 86f
left, 42f
perineural anesthesia, 121–24, 122f–124f
- Hip
caudal aspect of, 54
cranial aspect of, 53f, 54
deep dissection of, 53f
medial aspect of, 53–54, 55f
palpation, manipulation of, for lameness assessment, 104, 106f
- Hip joint (coxofemoral joint)
anatomy of, 54–56
anesthesia, 140, 140f
dislocation of, 380–82, 381f
infectious arthritis of, 383–84, 384f
OA of, 382–83, 382f–383f
physitis of, 383–84, 384f
- Hock. *See* Tarsus
- Hock joint. *See* Tarsal joint
- Hoof
corium relationship to, 7f
photomicrograph of, 9f
topography of, 6f
- Hoof crack, dorsal, 86f
- Hoof testers, 87f
- Hoof wall
anatomy of, 4–9, 6f–9f
coronary region of, three-dimensional dissection of, 8f
grooving of, 261f
- Horn, histological relationships of, 9f
- Hospital information system, 154
- Humeral-ulnar joint, caudolateral approach to, 135f
- Humerus
fractures of, diagnosis of, 330–31, 331f
osseus changes on, 334f
spiral nonarticular fracture of, 332f
stress fractures of, 161f, 331f
- Hyaluronan (HA)
benefits of, 416–17
common preparations of, 410t
corticosteroids, 409–10
intralesional, 411
intrasynovial, 409
parenteral, 406
- Iatrogenic tendon lesion, imaging of, 190f
- ICL. *See* Inferior carpal ligament
- Iliac wing fractures
clinical signs of, 390–91
diagnosis of, 391, 391f
treatment of, 391
- Image acquisition counts, 193t
- Imaging. *See also* Nuclear medicine; Ultrasonography
of canon bone, 173, 174f
CT benefits of, 216–17
DDFT, regions for, 182f
of DDFT, 172f, 174f, 176f–180f
of degenerative changes, 162f
detectors, 151
equipment, 149
for nuclear medicine, 193–94
film processing, 151

- Imaging. *See also* Nuclear medicine; Ultrasonography (*continued*)
of iatrogenic tendon lesion, 190f
of ligaments, 173, 174f
of osseous body, 162f
of osteochondrosis, 166f–167f
of osteomyelitis, 163f
of osteophytosis, 164f
protocol
for metacarpus, 171f
for ultrasound, 170–71, 171f
radiography, 149
of sclerosis, 162f
of SDFT, 172f, 174f, 176f–180f, 182f
of septic arthritis, 164f, 166f
of sesamoidean ligaments injury, 189f
of stifle, 165f–166f
of suspensory ligament injury, 188f
tendons, 173, 174f
transducer placement for, 183f–187f
viewing devices for, 151
x-ray machines for, 149–51
- Immobilization
of distal MC, 433, 435f
of elbow, 434, 435f
as initial treatment of, 432–33, 433t
by locations, 434f
of long bone fractures and luxations, 432–36, 433t, 434f–436f
methods of, 433–35, 434f–435f
of MT, 434
of phalanges, 433–34, 435f
of radius, 433–35, 435f
of tarsus, 435, 436f
of tibia, 435, 436f
- Impingement, of dorsal spinous process
clinical signs of, 395
generalizations of, 395
- Increased radiopharmaceutical uptake (IRU), 191f
- Infections
puncture wounds and, of calcaneal bursa, 430f
synovial, 436–38, 437f, 437t, 439t
- Infectious arthritis, of hip joint, 383–84, 384f
- Inferior carpal ligament (ICL)
palpation, manipulation of, for lameness assessment, 92
ultrasound of, 174f, 176f–177f
- Intersesamoidean desmitis, MRI diagnosis of, 214t
- Intertubercular bursa. *See* Bicipital bursitis
- Intra-articular carpal fractures
clinical signs of, 317–18
diagnosis of, 318, 318f
treatment of, 318–19, 318f–319f
- Intra-articular corticosteroids, 409t
- Intralesional medications
corticosteroids, 412–13
HA, 411
mesenchymal stem cell therapy, 412
PRP, 412
PSGAGs, 411–12
- Intrasynovial anesthesia
anesthetics, 124
general considerations, 124
guidelines for, 125, 125t
response to, 126
restraints used in, 126
skin preparation, 124
- Intrasynovial medications
ACS, 411
bone marrow mesenchymal stem cell therapy, 411
combination, with anesthetic, 411
corticosteroids, 408–9
fat-derived mesenchymal stem cell therapy, 411
HA, 409
Polyglycan®, 410
PSGAGs, 410
- Intravenous regional limb perfusions, using pneumatic tourniquet, 438f
- Inversion recovery (IR), 207
- IRU. *See* Increased radiopharmaceutical uptake
- Isoxsuprine, 406
- Joint anesthesia
coxofemoral, 140, 140f
DIP, 126–27, 127f–128f
DIT, 136–37, 137f
elbow, 132–34, 135f
FR, 139, 193f
LFT, 139–40, 140f
MCP/MTP fetlock, 128–30, 130f
MFT, 138–39, 138f
middle carpal joint, 131
proximal interphalangeal, 127–28, 129f
radiocarpal joint, 131–32
shoulder, 134–36, 135f
SI, 140–41, 141f
TC, 137–38
TMT, 136, 136f
- Joint disease, radiographic interpretation of, 163–65
types of, 165–67, 166f
- Jumping horses, occupation-related lameness in, 77t
- Keratoma
diagnosis of, 249–50, 250f
treatment of, 250
- Kimzey splint, 297f
- Lacerations
of DDFT, 441f
of MC/MT, 440, 440f
of MT, 440, 440f
- of SDFT, 441f
tendon, 438–39
clinical signs of, 439–40, 440f
diagnosis of, 440, 441f
flexor, 441f
treatment of, 440–41, 440f–441f
ultrasound assessment of, 182, 192f
- Lag screw, repair, of medial malleolus, 349f
- Lameness, occupation-related
horses with, 77t
types of, 76t
- Lameness, severe unilateral
causes, 429
clinical signs of, 429, 430f
treatment of, 429
- Lameness assessment
CT indications for, 219, 219f–221f
in forelimb, 84f
in hindlimb, 84f
history for, 83
MRI for, 216t
palpation, manipulation for
of back, 104, 107f–108f
of carpus, 93, 95f–96f
of femur, 103–4
for fetlock, 89–90, 89f–90f
of flexor tendons, 92, 95f
in foot, 85–87, 85f–87f
of forearm, 96–97, 97f
of hip, 104, 106f
of inferior carpal check ligament, 92
in MC/MT, 90, 91f–92f
of neck, 106
for pastern, 88, 88f–89f
of pelvis, 104, 106f
of shoulder, scapula, 97–98, 98f
of stifle, 101–3, 101f–104f
for suspensory ligament, 92, 94f
of tarsus, 98–100, 99f–100f
of tibia, 100–101, 101f–102f
signalment for, 83
visual exam for, at rest, 83–85
- Lameness diagnosis
for anatomic problem areas, 75
fundamentals of
classification of, 66
definition of, 65
in forelimb, 72–73, 73b, 73f
gaits, 67–69, 67f–69f
in hindlimb, 73–74, 73t
looking for, 66–67, 67f
stride, 69–72, 69f–72f
in hindlimb, 86f
for occupation-related problem areas, 75–76
- Lameness evaluation, of foals, 76–78, 78f
- Laminitis
example of, 259f
treatment of, 259–60, 261f
- Landing stride, 69–70, 71f

- Lateral aspect
 anatomy, nomenclature of, 20, 20f–21f
 of shoulder joint ligaments, 29f
 of tarsus, 36
 of thigh, 51–53, 52f–53f
- Lateral femorotibial joint (LFT), anesthesia, 139–40, 140f
- Lateral palmar block
 lateral approach, 119, 119f
 medial approach, 120, 120f
- Leg. *See* Crus
- Lesions
 diagnosis of, delayed phase for, 201f
 entosis-like, 214t
 iatrogenic tendon, imaging of, 190f
 MRI diagnosis of, 214t
 navicular disease and syndromes associated with, 227f
- LFT. *See* Lateral femorotibial joint
- Ligaments. *See also specific ligaments*
 of foot, 10f
 imaging of, 173, 174f
 of scapula, 28f–29f, 31, 36f
 signal intensity of, 208t
 supraspinous injuries to, 396–97, 397f
 ultrasonography assessment of, 169
 of pathology of, 175–78, 176f–177f
- Ligamentum nuchae, anatomy of, 61
- Loading stride, 70, 71f
- Local medications
 counterirritation from, 408
 DMSO, 407–8
 extracorporeal shock wave therapy, 408
 first aid, 407
 topical NSAIDs, 407
- Long bone fractures, luxations
 causes, 432
 definitive treatment of, 435–36
 diagnosis of, 432f
 first aid kit for, 433t
 immobilization treatment of, 432–36, 433t, 434f–436f
- Low palmar/four point block, 117, 118f
- Lumbar vertebrae, anatomy, 58f, 59
- Luxation and subluxation
 coxofemoral, 380–82, 381f
 long bone
 causes, 432
 definitive treatment of, 435–36
 diagnosis of, 432f
 first aid kit for, 433t
 immobilization treatment of, 432–36, 433t, 434f–436f
 of PIP, diagnosis of, 272f–274f, 273–74
 of SDFT, from calcaneus, 350–52, 351f–353f
 of tarsal joints, 349–50, 350f
- Magnetic resonance imaging (MRI)
 abnormalities of, 212, 212f–214f, 214t
 advantages of, 205
 artifacts for, 209–12, 211f
 case selection for, 205–6
 disadvantages of, 205
 equipment for, 203f–205f, 204–6
 images of, 203f–204f
 indications for, 205–6
 for lameness assessment, 216t
 of foot, 213t, 214f
 lesions identified with, 214t
 low-field, 205
 for MCP/MTP region, 216t
 of navicular disease, syndrome, 229f
 physics of, 202–4
 practicalities of, 204–6
 principles of, 202–4
 protocols for, 207–9, 208f, 209t–210t
 sequences for, 207–9, 208f
 signal intensity of, in different tissues, 208t
 ultrasound *vs.*, 189
- MC. *See* Metacarpus
- MCIII. *See* Metacarpus, dorsal
- MCP. *See* Metacarpal joint
- MCP/MTP fetlock. *See* Metacarpo/Metatarsophalangeal joint
- Meclofenamic acid (Arquel), 414–15
- Medial collateral ligament, 50f
- Medial cutaneous antebrachial blocks, 121, 121f
- Medial femorotibial joint (MFT)
 abnormalities of, 369f–370f
 anesthesia, 138–39, 138f
 synovitis and capsulitis of, OA of, 375
- Medial malleolus, lag screw repair of, 349f
- Meniscal injuries
 clinical signs of, 371–72, 372f
 diagnosis of, 372, 372f–373f
 tears, 373f
 treatment of, 372
 ultrasound of, 194f
- Mesenchymal stem cell therapy, intralesional, 412
- Metacarpal joint (MCP, osselets), traumatic OA of
 diagnosis of, 293, 294f
 treatment of, 293
- Metacarpo/Metatarsophalangeal joint (MCP/MTP fetlock)
 joint anesthesia for, 128–30, 130f
 MRI diagnosis of, 216t
- Metacarpus (MC)
 anatomy, nomenclature of, 14, 18
 distal, immobilization of, 433, 435f
 imaging protocol for, 171f
 MRI protocol for, 210t
 small
 exostosis, 304–6, 306f
 fractures of, 306–8, 307f–308f
- Metacarpus, dorsal (MCIII)
 complete fractures of, 303–4, 304f–305f
 condylar fractures of
 diagnosis of, 302–3, 302f
 treatment of, 303
 fracture of, 299–301, 300f–301f
 periostitis of, 299–301, 300f–301f
 stress fractures of, 299–301, 300f–301f
- Metacarpus/Metatarsus (MC/MT)
 lacerations of, 440, 440f
 lameness assessment of, palpation, manipulation for, 90, 91f–92f
 MRI protocol for, 210t
- Metaphyseal femoral fractures, clinical signs of, 377–78, 378f
- Metaphyseal tibial fractures
 clinical signs of, 359–60
 diagnosis of, 359f, 360
 treatment of, 360, 360f
- Metatarsus (MT)
 anatomy, nomenclature of, 32, 35f, 36
 fractures of, 306–8, 307f–308f
 immobilization of, 434
 lacerations of, 440, 440f
 medial dissection of, 38f
 MRI protocol for, 210t
- Metatarsus, dorsal (MTIII)
 complete fractures of, 303–4, 304f–305f
 condylar fractures of
 diagnosis of, 302–3, 302f
 treatment of, 303
- Methocarbamol. *See* Robaxin
- Methylsulfonylmethane (MSM), 417
- MFT. *See* Medial femorotibial joint
- MG/MT. *See* Metacarpus/Metatarsus
- Middle carpal joint
 anesthesia, 131
 approaches to, 134f
- Middle phalanx (P2), fractures
 clinical signs of, 275–77
 comminuted, 277, 278f
 diagnosis of, 276f, 277
 treatment of, 277, 277f–278f
- Midsagittal fractures, 238t
- Mineralization, of soft tissue, 157–58, 158f
- Minimum image acquisition counts, 193t
- MRI. *See* Magnetic resonance imaging
- MSM. *See* Methylsulfonylmethane
- MT. *See* Metatarsus
- Musculoskeletal system
 anatomy, functional of
 antebrachium, 22–23
 arm, shoulder, 25, 27f–28f
 carpus, 18, 20, 20f
 crus, 43, 44f–45f
 cubital joint, 23, 23f–24f, 25
 digit, fetlock, 3, 4f
 dorsal aspect, 18

- Musculoskeletal system (*continued*)
 extensor muscles, 20f, 23, 24f
 fetlock, 14, 15f–17f
 foot, 3
 hindlimb, 32, 35f, 36
 hip, 51–56, 52f–55f
 hoof wall, sole, frog, 4–9, 6f–9f
 lateral aspect, 20, 20f–21f
 medial aspect, 21, 21f
 metacarpus, 14, 18
 of neck, 61–62, 61f–62f
 nomenclature, usage, 3
 palmar aspect, 18, 19f, 21–22, 22f
 pastern, 11f–12f, 13–14
 positional, directional terms for, 4f
 stay apparatus, 31–32
 stifle, 43, 44f–49f, 46
 tarsal joint, 40–43, 41f–42f
 thigh, 51–56, 52f–55f
 of trunk, 61–62, 61f–62f
 examination of, 79–80
 of neck, 61–62, 61f–62f
 of scapula, 26f, 30–31
 scintigraphic examination of, imaging technique, 194–96, 196f
 of shoulder joint, 25, 28, 29f, 30
 of trunk, 61–62, 61f–62f
 ultrasound assessment of, 178–87, 191f–194f
- Myotomy, for fibrotic myopathy, 377f
- Naprosyn. *See* Naproxen
 Naproxen (Equiproxen®, Naprosyn), 415
- Natural balance shoes, 229f
- Navicular bone (distal sesamoid bone)
 anesthesia for, 143–44, 143f–144f
 attachments of, 13f
 views of, 13f
- Navicular disease and syndromes
 abnormalities in, 226t, 228f
 clinical signs of, 226–27
 fractures of, 230–31, 230f, 231f
 lesions associated with, 227f
 MRI of, 229f
 treatment of, 228–29
- Navicular wedge test, 113, 114f
- Neck
 muscles of, 61–62, 61f–62f
 palpation, manipulation of, for lameness assessment, 106
- Nerve blocks
 DBLPN, 123, 123f
 ulnar, 120, 121f
- Nerve injury. *See* Suprascapular nerve injury
- Nerves, of foot, 16f–17f
- Noncommittated fractures, of P1, treatment of, 279f, 281
- Nonseptic calcaneal bursitis, 353
- Nonsteroidal anti-inflammatories (NSAIDs)
 oral, 413–14, 413t
 parenteral, 405
 topical, 407
 types of, 413–15, 413t
- Nuchal bursitis, treatment of, 402
- Nuchal ligament desmopathy, treatment of, 402
- Nuclear medicine
 imaging equipment for, 193–94
 limitations of, 199–200
 principles of, 189–92
⁹⁹Tc in, 189–90, 195f
- Nuclear scintigraphy. *See also* Scintigraphic examination
 delayed phase of, 197
 of skeletal system, 196–97
 soft tissue phase of, 196–97, 197f
 vascular phase of, 196
- Nutraceuticals, GLN, 415
- OA. *See* Osteoarthritis
- Occupation-related lameness
 in draft horses, 77t
 in dressage horses, 77t
 in endurance horses, 77t
 in eventing horses, 77t
 in jumping horses, 77t
 in show or pleasure horses, 77t
 in standardbreds, 77t
 in thoroughbreds, 76t
 types of, 76t
- OCD. *See* Osteochondrosis
- Olecranon bursitis, 329–30, 329f
- Oral and nutritional medications, NSAID types of, 413–14, 413t
- Osselets. *See* Metacarpal joint
- Osseous body
 changes of, on humerus, 334f
 imaging of, 162f
- Ossification, of collateral cartilage, of distal phalanx, 241–43, 242f
- Osteoarthritis (OA). *See also* Traumatic OA
 of carpus
 clinical signs of, 320, 320f
 diagnosis of, 320, 321f
 treatment of, 320–21
 cervical facet joint, 402–3
 of coxofemoral joint, 382–83, 382f–383f
 of DIP, clinical signs of, 234–35, 234f
 distal tarsal, 343–46, 344f–345f
 facet joint, 399–401, 401f–402f
 MFT, 375
 MRI diagnosis of, 214t
 of PIP, 267–69, 268f–270f
- Osteochondral fractures (chip fractures)
 of carpal bones, 319f
 of P1, 286–87, 287f
- Osteochondral fragmentation
 MRI diagnosis of, 214t
 of P1, 288f
- Osteochondroma, of carpus, 322f
- Osteochondrosis (OCD)
 femoropatellar, 363–64, 365f
 imaging of, 166f–167f
 of PIP, 270–71, 271f
 of shoulder joint, 333–35, 335f
 of tarsocrural joint, 346–47, 346f–347f
- Osteomyelitis
 imaging of, 163f
 radiographic interpretation with, 161–63, 163f
- Osteophytosis, imaging of, 164f
- Osteostixis, 301f
- Overriding and impingement, of dorsal spinous process, generalizations of, 395
- P1. *See* Proximal phalanx
 P2. *See* Middle phalanx
 P3. *See* Distal phalanx
- Painful response, 108f
 test, for SI joint pain, 395f
- PAL. *See* Primary angular ligament
- Palmar anular desmitis, MRI diagnosis of, 214t
- Palmar anular ligament, drawing of, 12f
- Palmar aspect, anatomy, nomenclature of, 18, 19f, 21–22, 22f
- Palmar digital block (PD), 116, 116f
- Palmar process fractures, diagnosis of, delayed phase for, 200f
- Palpation, manipulation
 of bones, 93f
 of collateral ligaments, 89f
 of cubital joint, 97f
 of digital flexor synovial sheath, 90f
 of ECR, 95f
 of extensor carpi radialis, 95f
 for lameness assessment
 of back, 104, 107f–108f
 of carpus, 93, 95f–96f
 of collateral ligaments, 89f
 of femur, 103–4
 for fetlock, 89–90, 89f–90f
 of flexor tendons, 92, 95f
 in foot, 85–87, 85f–87f
 of forearm, 96–97, 97f
 of hip, 104, 106f
 of inferior carpal check ligament, 92
 for MC, MT, 90, 91f–92f
 of neck, 106
 for pastern, 88, 88f–89f
 of pelvis, 104, 106f
 of sesamoidean ligaments, 88f
 of shoulder, scapula, 97–98, 98f
 of stifle, 101–3, 101f–104f
 for suspensory ligament, 92, 94f
 of tarsus, 98–100, 99f–100f
 of tibia, 100–101, 101f–102f
 of metacarpal bones, 93f

- of SDFT, 100f–101f
- Parenteral medications
 - bisphosphonates, 406
 - HA, 406
 - isoxsuprine, 406
 - NSAIDs, 405
 - PSGAGs, 405–6
 - Robaxin, 407
 - tetracyclines, 407
- Pastern
 - anatomy, nomenclature of, 11f–12f, 13–14
 - examination of, 187f
 - injuries
 - DDFT, 284–86, 285f–286f
 - SDFT, 284–86, 285f
 - lameness assessment in, palpation, manipulation for, 88, 88f–89f
- Patella
 - fractures of, 364–67, 366f
 - upward fixation of, 367–68, 367f, 381f
- PBZ. *See* Phenylbutazone
- PD. *See* Palmar digital block
- Pedal osteitis (PO)
 - diagnosis of, 239, 240f
 - treatment of, 239–40
- Pelvic fractures
 - clinical signs of, 389, 390f
 - diagnosis of, 389, 390f
 - treatment of, 390
- Pelvic limb
 - bones of, 34f
 - stay apparatus of, 56, 57f
- Pelvis
 - anatomy of, 56
 - palpation, manipulation of, for lameness assessment, 104, 106f
- Penetrating injuries, of foot
 - diagnosis of, 248–49
 - treatment of, 249
- Perineural anesthesia
 - anesthetics, 114
 - assessment in, for block responses, 115–16
 - general considerations, 114
 - guidelines for, 115t
 - restraints used in, 114–15
 - skin preparation, 114
- Perineural blocks
 - forelimb
 - abaxial sesamoid block, 117, 117f
 - forelimb, 116–24, 116f–124f
 - high palmar/high four-point block, 118, 119f
 - high PD block, 116–17, 116f
 - high two-point block, 120, 120f
 - hindlimb, 121–24, 122f–124f
 - lateral palmar block, 119, 119f, 120, 120f
 - low palmar/four point block, 117, 118f
 - medial cutaneous antebrachial blocks, 121, 121f
 - PD block, 116, 116f
 - ulnar nerve block, 120, 121f
 - hindlimb
 - DBLPN, 123, 123f
 - high plantar/sub-tarsal blocks, 122–23, 122f
 - low plantar/six-point block, 122, 122f
 - peroneal, 123–24, 123f–124f
 - tibial blocks, 123–24, 123f–124f
 - responses, 115
 - types of, 115t
- Periosteum
 - histological relationship to, 9f
 - reactions of, 159f, 160
- Periostitis, of dorsal MC, 299–301, 300f–301f
- Peroneal blocks, 123–24, 123f–124f
- Peroneus tertius
 - clinical signs of, 356–57
 - treatment of, 357
- Phalanges, immobilization of, 433–34, 435f
- Phenylbutazone (PBZ; Butazolidin®, Butatron™, Bizolin®, Phenylbute™, Phenylzone®, Equiphen®, Butequine®, Superiorbute®, Equizone 100™), 414
- Phenylbute™. *See* Phenylbutazone
- Phenylzone®. *See* Phenylbutazone
- γ-photos, 195f
- Photomultiplier tubes, 195f
- Physitis, of hip joint, 383–84, 384f
- Picture archive and communication system, 154
- PIP. *See* Proximal interphalangeal joint
- Plantar aspect, of tarsus, 39–40
- Platelet-rich plasma (PRP), intralesional, 412
- Pneumatic tourniquet, intravenous regional limb perfusions using, 438f
- PO. *See* Pedal osteitis
- Podotrochlear apparatus, injuries to, 232f
 - treatment of, 232
- Point of break-over, in trimming, 422f
- Polyglycan®, 410
- Polysulfated glycosaminoglycans (PSGAGs)
 - intralesional, 411–12
 - intrasynovial, 410
 - parenteral, 405
- Prepurchase examination
 - performing of, 78–81, 81t
 - reporting results of, 80–81, 81t
- Primary angular ligament (PAL), ultrasound assessment of, 180f
- Proximal digital annular ligament, drawing of, 12f
- Proximal interphalangeal joint (PIP)
 - anesthesia, 127–28, 129f
 - luxation and subluxation of, diagnosis of, 272f–274f, 273–74
 - OA of, 267–69, 268f–270f
 - OC of, 270–71, 271f
- Proximal phalanx (P1)
 - fractures of
 - of comminuted fractures, 282, 282f
 - noncomminuted, 279f, 281
 - osteocondral fractures of
 - diagnosis of, 287, 287f
 - treatment of, 287
 - osteocondral fragmentation of, 288f
- Proximal radius, immobilization of, 433–35, 435f
- Proximal sesamoid bones, fractures of, treatment of, 288f, 290f–291f, 291
- Proximal suspensory desmitis, 217f
- Proximal tibial physis, fractures of, 362–63, 362f–363f
- PRP. *See* Platelet-rich plasma
- PSGAGs. *See* Polysulfated glycosaminoglycans
- Punctures
 - infection and, 430f
 - ultrasound assessment of, 182, 192f
- PVC splint, 435f
- Quadrisol®. *See* Vedaprofen
- Quarter cracks
 - clinical signs of, 255–56
 - treatment of, 256, 256f–257f
 - wire placement for, 257f
- Radiation safety, 155–56
- Radiocarpal joint
 - anesthesia, 131–32
 - dorsal flexed, 134f
- Radiofrequency coils, 204f
- Radiographic interpretation. *See also* Ultrasonography
 - of bones, 159–63, 160f–163f
 - of joint conditions, 165–67, 166f
 - limitations of, 167–68
 - with osteomyelitis, 161–63, 163f
 - principles of, 156–57
 - of soft tissue, 157, 157f
 - of synovial joints, 163
 - in joint disease, 163–65
 - ultrasonography *vs.*, 191f
- Radius
 - fractures of
 - diagnosis of, 323–24, 324f
 - treatment of, 324, 325f
 - immobilization of, 433–35, 435f
 - proximal, immobilization of, 433–35, 435f
- Reciprocal apparatus, lateral view of, 42f

- Repetition time (TR), 207
 Rhabdomyolysis, delayed phase of, 202f
 Rimadyl®. *See* Carprofen
 Robaxin (Methocarbamol), 407
- Sacral ligament, schematic drawing of, 394f
 Sacroiliac joint (SI)
 anatomy of, 60–61, 61f
 anesthesia, 140–41, 141f
 diseases of
 clinical signs of, 394, 394f
 generalizations of, 393–94, 394f
 treatment of, 395
 injection of, 141f
 minimum image acquisition counts in, 193t
 pain provocation test for, 395f
 schematic drawing of, 394f
 Sacrum
 anatomy, 58f, 59
 fractures of, 392–93, 393f
 Sagittal fractures, 347–48, 348f
 of central tarsal bones, 347–48, 348f
 Scapula
 ligaments of, 28f–29f, 31, 36f
 muscles overlying, 26f, 30–31
 palpation, manipulation of, for lameness assessment, 97–98, 98f
 Scintigraphic examination. *See also* Nuclear scintigraphy
 of musculoskeletal system, imaging technique, 194–96, 196f
 Scintigraphic signs, of disease
 delayed phase, 197, 198f–202f, 199
 soft tissue phase, 197, 197f–198f
 Sclerosis, imaging of, 162f
 SCLs. *See* Subchondral cystic lesions
 SDFT. *See* Superficial digital flexor ligament
 SE. *See* Spin echo
 Septic arthritis, imaging of, 164f, 166f
 Septic calcaneal bursitis, 353
 Sesamoid block, abaxial, 117, 117f
 Sesamoidean ligaments
 drawing of, 12f
 injury of, imaging of, 189f
 palpation of, 88f
 ultrasound of, 174f, 176f
 Sesamoid fracture
 abaxial, 288f–289f
 apical, 288f, 290f–291f
 Sesamoiditis
 diagnosis of, 292, 292f
 treatment of, 292
 Sheared heels, 252f–253f
 Shin splints, 299–301, 300f–301f
 Shoes
 center of pressure for, 420–21, 420f–422f
 corrective, 417–19, 418f, 419t
 distribution of force by, 421–22
 for ease of movement, 421, 422f
 foot motion and, during flight, 422–23
 full bar, for distal phalanx fracture, 238f
 natural balance, 229f
 for protection, 423
 shock absorption of, 419–20
 straight-bar, 256f
 for support, 423
 weight-bearing, 420–21
 Shoulder
 anatomy, nomenclature of, 25, 27f–28f
 medial view of, 26f
 minimum image acquisition counts in, 193t
 Shoulder joint
 anesthesia, 134–36, 135f
 craniolateral, lateral approaches to, 135f
 draining tract at, 193f
 flexion tests, manipulation, upper forelimb and, 110–11, 111f
 ligaments
 flexor muscles related to, 30
 lateral aspect of, 29f
 medial view of, 26f
 muscles substituting for, 25, 28, 29f, 30
 right, 28f
 OCD of, treatment of, 334, 335f
 palpation, manipulation of, for lameness assessment, 97–98, 98f
 Show and pleasure horses, occupation-related lameness in, 77t
 SI. *See* Sacroiliac joint
 Sidebone. *See* Distal phalanx
 Siemens Symphony 1.5 Tesla
 image of, 203f
 protocols for, 209t–210t
 Skeletal system, nuclear scintigraphy of, 196–97
 Slab fractures, of central tarsal bones, 347–48, 348f
 Small MC bone exostosis
 clinical signs of, 305
 diagnosis of, 305–6, 306f
 treatment of, 306, 306f
 Soft tissue
 bruising of, 430f
 entheses in, 159, 159f
 gas in, 158–59, 159f
 mineralization of, 157–58, 158f
 radiographic interpretation of, 157, 157f
 thickening of, 157, 157f, 160f
 Soft tissue phase, of nuclear scintigraphy, 196–97, 197f–198f
 Sole
 anatomy of, 4–9, 6f–9f
 histological relationships in, 9f
 topography of, 6f
 Sole bruises, 244f
 clinical signs of, 243
 treatment of, 243–44
 Spavin test, 111, 112f
 Spine, minimum image acquisition counts in, 193t
 Spin echo (SE), 207
 Spinous process
 fractures of, 397
 overriding and impingement of
 clinical signs of, 395
 generalizations of, 395
 Spiral nonarticular fracture, of humerus, 332f
 Splints
 caudal, 435f
 diagnosis of, 305–6, 306f
 Kimzey, 297f
 PVC, 435f
 in radial fracture, 325f
 shin, 299–301, 300f–301f
 treatment of, 306, 306f
 Spondylosis
 diagnosis of, 399, 399f
 treatment of, 399
 Stance, 70–71, 71f
 Standardbreds, occupation-related lameness in, 77t
 Stay apparatus
 of pelvic limb, 56, 57f
 of thoracic limb, 31–32
 Stem cell therapy, 411–12
 Stifle (genu)
 anatomy, nomenclature of, 43, 44f–49f, 46
 caudal aspect of, 48f
 deep dissection of, 49f
 deep dissection of, 47f
 dorsal view of, 44f
 flexion test of, 112, 113f
 imaging of, 165f–166f
 medial aspect of, deep dissection of, 50f
 minimum image acquisition counts in, 193t
 palpation, manipulation of, for lameness assessment, 101–3, 101f–104f
 SCLs of, 368–70, 369f
 superficial dissection of, 45f
 Stifle joint
 anatomy, nomenclature of, 47–51
 femoropatellar joint, 47–49
 femorotibial joint, 49–51, 49f
 movements of, 51
 Straight-bar shoes, 256f
 Stress fractures
 delayed phase diagnosis of, 199f
 of dorsal MC, 299–301, 300f–301f
 humeral, 161f, 331f
 tibial, 161f
 diagnosis of, 358, 359f
 treatment of, 358–59

- Stride
 landing, 69–70, 71f
 length of, 72, 72f
 loading, 70, 71f
 phases of, 69–72, 69f–72f
 swing phase of, 72, 72f
- Stringhalt, treatment of, 358, 358f
- Subchondral bone injury, MRI
 diagnosis of, 214t
- Subchondral cystic lesions (SCLs). *See* *also* Fetlock subchondral cystic lesions
 of distal phalanx
 diagnosis of, 241f
 treatment of, 241
 radiograph of, 371f
 of stifle, 368–70, 369f
- Subluxation. *See* Luxation and subluxation
- Superficial digital flexor ligament (SDFT)
 drawing of, 12f
 imaging of, 172f, 174f, 176f–180f
 regions for, 182f
 lacerations of, 441f
 luxation of, from calcaneus, 350–52, 351f–353f
 palmaromedial view of, with flexor retinaculum cut, 22f
 palpation of, 100f–101f
 pastern injuries, treatment of, 285–86
 tendonitis
 diagnosis of, 313, 314f
 MRI diagnosis of, 214t
 treatment of, 313–14
 transducers in, 183f–187f
 ultrasound assessment of, 172f, 174f, 176f–180f
- Superiorbute®. *See* Phenylbutazone
- Supraglenoid tubercle (tuberosity), fractures of, 337–38, 337f
- Suprascapular nerve injury (SWEENY)
 clinical signs of, 336, 336f
 treatment of, 336–37
- Supraspinous ligament injuries
 diagnosis of, 396, 397f
 treatment of, 396
- Suspensory apparatus, traumatic rupture of
 diagnosis of, 296, 296f
 treatment of, 296, 297f
- Suspensory ligament
 attachments of, 309f
 branch, ultrasound assessment of, 181f
 desmitis, 310f
 degenerative, 311–14, 312f–314f
 diagnosis of, 310, 311f
 treatment of, 310
 injury, imaging of, 188f
 palpation, manipulation of, for lameness assessment, 92, 94f
- Sustentaculum tali, fractures of, 355f
- SWEENY. *See* Suprascapular nerve injury
- Swing phase, 72, 72f
- Swollen limb, severe
 causes, 431
 clinical signs of, 431, 431f
 treatment of, 431
- Symmetrical distal displacement, 258f
- Synovial infections
 antimicrobials for, 437t, 439t
 causes, 436
 diagnosis of, 436–37, 437f
 treatment of, 437, 437t, 439t
- Synovial joints, radiographic interpretation of, 163
 in joint disease, 163–65
- Synovial sheaths, ultrasound of, 178–80, 191f
- Synovitis, MFT, 375
- Systemic and parenteral medications
 bisphosphonates, 406
 HA, 406
 isoxsuprine, 406
 NSAIDs, 405
 PSGAGs, 405–6
 Robaxin, 407
 tetracyclines, 407
- Tarsal bones
 fractures of
 delayed phase diagnosing of, 200f
 SLAB/sagittal, 347–48, 348f
 third, SLAB/sagittal fractures of, 347–48, 348f
- Tarsal joint (hock joint)
 anatomy, nomenclature of, 40–43, 41f–42f
 luxation and subluxation of, 349–50, 350f
- Tarsal sheath anesthesia, 143
- Tarsal sheath tenosynovitis (thoroughpin), 354–56, 354f–355f
- Tarsocrural joint (TC)
 anesthesia, 137–38
 dorsomedial, dorsolateral approach to, 137f
 movements of, 35f, 37f, 41, 43
 OCD of, 346–47, 346f–347f
- Tarsometatarsal joint (TMT)
 anesthesia, 136, 136f
 lateral approach to, 136f
- Tarsus (hock)
 anatomy, nomenclature of, 34f, 36
 capped hock, 352–53
 dorsal aspect, 36, 37f
 dorsal dissection of, 37f
 dorsal view of, 41f, 44f
 flexion tests, manipulation of, 111, 112f
 immobilization of, 435, 436f
 lateral aspect of, 36
 medial aspect of, 36, 38f–39f, 39
 medial dissection of, 38f
 medial view of, 39f
 minimum image acquisition counts in, 193t
 MRI protocol for, 210t
 palpation, manipulation of, for lameness assessment, 98–100, 99f–100f
 plantar aspect of, 39–40
 superficial dissection of, 45f
- TC. *See* Tarsocrural joint
- TE. *See* Echo time
- Technetium-99m (^{99m}Tc), in nuclear medicine, 189–90, 195f
- Tendonitis
 DDFT, 214t
 SDFT, 313–14, 314f
 MRI diagnosis of, 214t
 ultrasound of, 177–78
- Tendon lacerations
 clinical signs of, 439–40, 440f
 diagnosis of, 440, 441f
 flexor, 441f
 treatment of, 440–41, 440f–441f
- Tendons. *See also specific tendons*
 imaging of, 173, 174f
 signal intensity of, 208t
 ultrasonography assessment of, 169, 175–78, 176f–177f
- Tendon sheath anesthesia
 carpal sheath, 142–43
 DFTS, 141–42, 141f–142f
 ECR, 143
 tarsal, 143
- Tenosynovitis
 carpal sheath, 321–23, 322f–323f
 of DFTS
 clinical signs of, 298, 298f
 diagnosis of, 298–99, 298f
 treatment of, 299, 299f
 tarsal sheath, 354–56, 354f–355f
 ultrasonography of, 191f
- Tenotomy, of DDFT, 262f
- Tetracyclines, 407
- Thickening, soft tissue, 157, 157f, 160f
- Thigh
 atrophy of, 84f
 caudal aspect of, 54
 cranial aspect of, 53f, 54
 lateral aspect of, 51–53, 52f–53f
 medial aspect of, 53–54, 55f
- Third tarsal bones, SLAB/sagittal fractures of, 347–48, 348f
- Thoracic limb
 anatomy, nomenclature of, 3, 4f
 bones of, 5f
 stay apparatus of, 31–32
- Thoracic vertebrae, anatomy, 58f, 59
- Thoroughbreds, occupation-related lameness in, 76t
- Thoroughpin. *See* Tarsal sheath tenosynovitis
- Thrush
 clinical signs of, 245–46
 treatment of, 246

- Tibia
 diaphyseal, metaphyseal fractures
 of, 359–60, 359f–360f
 immobilization of, 435, 436f
 palpation, manipulation of, for
 lameness assessment, 100–101,
 101f–102f
 Tibial blocks, 123–24, 123f–124f
 Tibial malleoli, fractures of, 348–49
 Tibial physis, proximal, fractures of,
 362–63, 362f–363f
 Tibial stress fractures, 161f
 diagnosis of, 358, 359f
 treatment of, 358–59
 Tibial tuberosity
 diagnosis of, 361–62
 treatment of, 361–62, 361f
 TMT. *See* Tarsometatarsal joint
 Toe cracks
 clinical signs of, 255–56
 treatment of, 256, 256f–257f
 Toe-out, hock-in stance, 106f
 Topical, local medications
 counterirritation from, 408
 DMSO, 407–8
 extracorporeal shock wave therapy,
 408
 first aid, 407
 topical NSAIDs, 407
 TR. *See* Repetition time
 Transducers
 in DDFT, 183f–187f
 for imaging, 183f–187f
 in SDFT, 183f–187f
 Traumatic OA, of MCP
 diagnosis of, 293, 294f
 treatment of, 293
 Trochanteric bursa anesthesia, 145,
 146f
 Trot, image of, 68f
 Trunk, muscles of, 61–62, 61f–62f
 Tuber coxae fractures
 clinical signs of, 391, 392f
 diagnosis of, 391, 392f
 treatment of, 391–92
 Tuberosity
 supraglenoid tubercle, 337–38,
 337f
 tibial
 diagnosis of, 361–62
 treatment of, 361–62, 361f
 Type II distal femoral fractures,
 379f
 Type IV fracture, 240f
 UFP. *See* Upward fixation of patella
 Ulna, fractures of
 clinical signs of, 327, 327f
 diagnosis of, 327, 327f
 treatment of, 327, 327f–328f
 Ulnaris lateralis. *See* Extensor carpi
 ulnaris
 Ulnar nerve block, 120, 121f
 Ultrasonography
 assessment with
 of bones, 180, 182, 191f
 of DDFT, 172f, 174f, 176f–180f,
 286f
 of foreign bodies, 182, 185, 193f
 ICL of, 174f, 176f–177f
 imaging protocol for, 170–71,
 171f
 of joints, 187, 194f
 of lacerations, 182, 192f
 limitations of, 187–89
 of meniscus damage, 194f
 MRI *vs.*, 189
 of musculoskeletal system,
 178–87, 191f–194f
 of PAL, 180f
 patient preparation for, 170–75,
 171f–174f
 of punctures, 182, 192f
 radiographs *vs.*, 191f
 science of, 168–69
 of SDFT, 172f, 174f, 176f–180f
 of sesamoidean ligaments, 174f,
 176f
 of suspensory ligament branch,
 181f
 of synovial sheaths, 178–80, 191f
 of tendinitis, 177–78
 of tendons, 169, 175–78,
 176f–177f
 of tenosynovitis, 191f
 for ligament pathology, 175–78,
 176f–177f
 of ligaments, 169
 Upward fixation of patella (UFP)
 clinical signs of, 368
 hip dislocation and, 381f
 treatment of, 368
 Vascular phase, of nuclear
 scintigraphy, 196
 Vedaprofen (Quadrisol®), 415
 Vertebral articulations, 60
 Vertebral column
 axial components of, 56, 58f
 caudal vertebrae, 60
 cervical vertebrae, 58–59, 58f
 lumbar vertebrae, 58f, 59
 thoracic vertebrae, 58f, 59
 Vertebral facet joint syndrome,
 399–401, 401f–402f
 Vertebral fractures
 clinical signs of, 397–98
 treatment of, 398
 Viewing devices, for imaging, 151
 Visual examination
 at exercise, 107–8
 for lameness assessment, at rest,
 83–85
 Walk, image of, 67f
 Weight-bearing shoes, 420–21
 White line disease
 clinical signs of, 246–47
 treatment of, 247
 Wires, for quarter crack repair, 257f
 X-ray machines, for imaging,
 149–51
 Zenecarp. *See* Carprofen