

Index

- abdominal palpation per rectum
 - abnormal findings, 26–7
 - cecum, 28, 30–31
 - kidney and spleen, 35
 - large colon distention, 28–9, 32–3
 - peritoneal cavity, 35–6
 - rectal tear, 33–4
 - small (descending) colon, 33
 - small intestine, 27–9
 - urogenital tract, 34–5
 - complications, 36–7
 - diagnoses, 28
 - differential diagnoses, 27
 - indications, 22–3
 - indications for referral, 74
 - mild colic, 46
 - normal anatomy
 - abdominal cavity, 25
 - bladder, 24
 - cecum, 25
 - inguinal rings, 25
 - kidney and spleen, 25
 - large colon, 26
 - pelvic region, 24
 - rectum, 24
 - small (descending) colon, 25–6
 - uterus and ovaries, 25
 - preparation
 - lubrication, 23
 - materials required, 23
 - rectal relaxation, 23–4
 - rectal sleeve, 23
 - restraint, 23
 - sedation, 24
 - procedure, 24–36
- abdominal radiography
 - diaphragmatic hernia, 157
 - enterolithiasis, 155–6
 - equipment
 - cassettes/detector and holder, 150–152
 - radiolucent markers, 151
 - safety equipment, 151
 - X-ray generator, 150
 - foals, 157–9
 - gut fill, 152
 - indications, 149–50
 - intestinal sand accumulation, 156–7
 - metallic foreign material, 157
 - sedation/restraint, 152
- abdominal sonographic evaluation
 - gastric distension and impaction, 131–4
 - indications, 117
 - intussusceptions, 127
 - large colon displacements

- abdominal sonographic evaluation (*cont'd*)
 - large colon volvulus, 122–3
 - left dorsal displacement, 121–2
 - right dorsal displacement, 120–121
- large intestinal impactions
 - large colon or cecal impaction, 123–4
 - meconium retention, 125
 - sand impactions, 124
 - small colon impactions, 124
- patient preparation, 117
- procedure
 - abdominal examination (complete), 118–20
 - FLASH, 118
 - transrectal examination, 120
- right dorsal colitis, 125–6
- small intestinal lesions
 - adhesions, 130–131, 134
 - idiopathic small intestinal muscular hypertrophy, 130, 133
 - strangulating *vs.* nonstrangulating obstruction, 128–30
- transducer selection, 117
- abdominocentesis
 - complications, 89–90
 - indications, 87–8
 - preparation, 88
 - procedure, 88–9
- acid-base imbalances, 80–81
- adhesions
 - long-term recovery, 294
 - miniature horses, 289
 - postoperative complication, 255–6
 - previous colic surgery, 9
 - sonographic evaluation, 130–131
- adult horses
 - abdominal radiography
 - caudodorsal radiograph, 153–4
 - caudoventral radiograph, 153, 156
 - cranioventral radiograph, 153, 155
 - metallic foreign objects, 157
 - mid-abdominal radiograph, 153, 155
 - antimicrobial drug concentrations, 339
 - arterial blood pressure, 102
 - biochemistry in, 338
 - colloid oncotic pressure, 100
 - drug dosages, 329–35
 - energy and protein requirements for, 302, 308
 - enteral nutrition, 303
 - fecal output, 7
 - glucose homeostatic dysfunction, 83
 - heart rate, 16
 - hematology in, 337
 - IV catheterization, 104, 107
 - IV fluid therapy
 - electrolyte replacement, 110
 - maintenance fluid rate, 110
 - replacement/resuscitation fluid, 109
 - colloids, 112–14
 - Landis-Pappenheimer equation for, 100
 - mild colic, 46–8
 - parenteral nutrition formulations for, 309
 - physical examination, 14–21
 - postoperative patient care
 - central venous pressure, 234
 - fluid therapy plan, 237
 - mean arterial pressure, 235
 - patient assessment, 231
 - physical examination, 232
 - urine-specific gravity, 234
 - sonographic examination, 118–20
 - urine output, 102
 - water consumption, 7
- alkaline phosphatase (ALP), 82
- alpha-2 adrenergic agonists, 17, 239
 - adverse effects and contraindications, 55
 - alpha-2 antagonists, 55
 - clinical use, 54–5
 - dose rates, 330, 335
 - mechanism of action, 54
- aminoglycosides
 - dose rates, 329, 332
 - toxicity, 9
 - for young foals and weanlings, 281
- analgesia
 - alpha-2 adrenergic agonists
 - adverse effects and contraindications, 55
 - alpha-2 antagonists, 55
 - clinical use, 54–5
 - dose rates, 330, 335
 - mechanism of action, 54
 - hyoscine-N-butylbromide
 - adverse effects and contraindications, 59
 - clinical use, 58–9
 - dose rate, 330
 - mechanism of action, 58
 - indications, 51–2
 - intravenous lidocaine
 - adverse effects and contraindications, 58
 - clinical use, 57–8
 - complications, 58

- dose rate, 332
- mechanism of action, 57
- mild colic, 46–7
- nonsteroidal anti-inflammatory drugs
 - adverse effects and contraindications, 53
 - clinical use, 52–3
 - dose rate, 331–4
 - mechanism of action, 52
 - mild colic, 46–7
- opioids
 - adverse effects and contraindications, 56–7
 - antagonists, 56
 - clinical use, 56
 - dose rate, 329
 - mechanism of action, 55
 - postoperative use, 238–40, 249–50
- Anoplocephala perfoliata* (tapeworms)
 - definition, 318
 - diagnosis and management, 319–20
 - life cycle, 318
 - prevention of colic, 297
 - pathology and clinical disease, 318
 - cecal perforation, 215
 - cecocolic intussusception, 215
 - ileal impaction, 208
 - ileocecal intussusception, 213
- anthelmintics, 319
 - gastrointestinal parasite control, 8–9
 - Anoplocephala perfoliata*, 318
 - ascarid impaction, 322
 - cyathostomes, 320–321
 - fecal egg count reduction test, 317
 - pasture management, 317
 - Strongylus vulgaris*, 323
 - resistance, 297, 317
- arrhythmia, 17
- arterial oxygen content (CaO₂), 101
- ascariasis, 321–2
 - impaction, 208–9
 - prevention, 209, 322
 - prevention of colic, 297
 - treatment, 209
 - young foals and weanlings, 280
- ascarid impaction, 208–209
- aspartate transaminase (AST), 82
- benzimidazole anthelmintics, 209, 321, 322
- biosecurity program
 - vs. biocontainment, 263
 - definition and terminology, 263
- for equine colic patient at home
 - cleaning and disinfection, 271–2
 - general recommendations, 270–271
 - MRSA, 273
 - shedding *Salmonella*, 272–3
- for hospitalized equine colic patient
 - monitoring and surveillance, 269–70
 - preventive measures, 269
 - shedding *Salmonella*, 268
- hospital acquired infections, 263–4
- infection control programs, 264
- monitoring and surveillance
 - hospital environment, 268
 - patient, 267–8
- preventive measures
 - antimicrobial use, 267
 - barrier precautions and protective clothing, 266
 - Biosecurity Officer, 267
 - cleaning and disinfection, 265–6
 - hand hygiene, 266
 - high and low risk patients, 265
 - training and education, 267
 - waste disposal, 266–7
- broad ligament hematoma
 - abdominal palpation, 27, 34–5
 - cause of colic and treatment, 282–4
 - sonographic examination, 120, 136
- broodmares
 - postpartum mares, 284–5
 - pregnant mares, 284
- Buscopan®
 - abdominal palpation per rectum, 23, 24
 - adverse effects and contraindications, 59
 - clinical use, 58–9
 - dose rate, 330
 - mechanism of action, 58
 - mild colic, 47
- butorphanol *see also* opioids
 - dose rate, 329
 - for young foals and weanlings, 281
 - mild colic, 46–7
 - postoperative pain, 250
 - postoperative patient care, 239
 - trocharization, 161
- cardiac output (Q), 101
- cecal impaction, 9, 10
 - abdominal palpation per rectum, 28, 30–31
 - clinical signs, 214
 - medical treatment, 214

- cecal impaction (*cont'd*)
 - necropsy, 215
 - sonography, 123–4
 - surgical treatment, 168, 214
 - typhlotomy, 190
- cecal perforation, 168, 215
- cecal tympany, 28, 30–31
- cecocecal intussusception, 214–15 *see also* cecocolic intussusception
- cecocolic intussusception, 198, 214
 - abdominal palpation per rectum, 28, 30–31
 - clinical signs, 215
 - sonographic evaluation, 127, 130
 - surgical treatment, 215
- cecum
 - abdominal palpation per rectum, 28, 30–31
 - celiotomy, 178–9
- ceftiofur, 330
- colic surgery, 175
 - for young foals and weanlings, 281
 - postoperative patient care, 240
- celiotomy *see also* colic surgery
 - abdominal organs, 181–2
 - approaches
 - ventral, 177
 - lateral, 185
 - cecum, 178–9
 - indications (general), 164–6
 - large colon, 182–3
 - small colon, 182
 - small intestine, 179–81
 - transverse colon, 182
- central venous pressure (CVP), 102
- clinical laboratory data
 - clinical chemistry
 - acid-base evaluation, 80–81
 - electrolyte imbalance, 81–2
 - hepatobiliary assessment, 82–3
 - metabolic indicators, 83
 - renal parameters, 83
 - coagulation
 - inhibitors, 84–5
 - platelets, 83–4
 - secondary hemostasis, 84
 - hematology
 - erythrogram, 79
 - leukogram, 79–80
 - indications, 78–9
- coagulation
 - inhibitors, 84–5
 - platelets, 83–4
- secondary hemostasis, 84
- colic
 - donkeys
 - causes, 285–6
 - hyperlipemia, 288–9
 - drug dosages, 329–35
 - geriatric horses
 - causes, 286
 - management and complications, 287
 - mild (*see* mild colic)
 - miniature horses
 - causes, 287
 - management and complications, 287–8
 - ponies
 - causes, 287
 - management and complications, 287–8
 - postpartum mares, 284–5
 - pregnant mares, 282–4
 - risk factors and prevention
 - colonic microbiota, 297
 - dental care, 298
 - diet, 296–7
 - exercise regimen, 297
 - gastrointestinal parasites, 297
 - genetic predisposition, 295
 - recurrent colic, 295
 - risk factors, 296–7
 - seasonal pattern, 296
 - transportation, 297
 - water deprivation, 297
- colic surgery
 - anastomosis, 192
 - end-to-end technique, 195–7
 - ileocecostomy, 194–5
 - jejunocecostomy, 193–4
 - jejunoileostomy, 193
 - jejunojejunostomy, 193
 - side-to-side anastomosis, 197
 - small colon resection, 197–202
 - colopexy, 202
 - complications
 - diarrhea, 248–9
 - endotoxemia and the associated SIRS, 246–8
 - fever, 244–6
 - incisional complications, 256–8
 - ischemia-reperfusion injury, 246
 - postoperative ileus, 251–3
 - postoperative intraperitoneal adhesions, 255–6
 - recurrent colic, 249–51
 - septic peritonitis, 253–4

- diagnosis and correction, 198–201
- enterotomy, 190–191
- indications, 164–6, 174
- laparoscopy, 177, 187–8
- lateral approaches, 196–7
 - body wall closure, 186–7
 - celiotomy, 186
 - incision, 185–6
- long-term recovery
 - complications, 294
 - feeding, 294
 - incisional care, 293–5
 - postoperative management, 292–4
 - return to use, 295
 - survival, 294–5
- needle decompression, 189
- partial typhlectomy, 191–2
- surgical preparation
 - analgesic drugs, 174–5
 - anesthesia, 175
 - antimicrobial drugs, 175
 - cardiovascular stabilization, 174
 - surgical site preparation, 176
- ventral approaches, 176
 - body wall closure, 183–5
 - celiotomy (*see* celiotomy)
 - incision, 177–8
- colitis, 9, 10, 169, 215–6 *see also* diarrhea
- colopexy, 202
- COX *see* cyclooxygenase (COX)
- CVP *see* central venous pressure (CVP)
- cyathostomes
 - clinical disease, 320
 - definition, 320
 - diagnosis and management, 320–321
 - life cycle, 320
 - pathology, 320
 - prevention of colic, 297
- cyclooxygenase (COX), 46
 - ischemia-reperfusion injury, 246
 - non-steroidal anti-inflammatory drugs, 46, 52–4
 - right dorsal colitis, 216
- detomidine, 52, 54–5, 74, 165, 239 *see also* alpha-2 agonists
 - abdominal palpation per rectum, 74
 - causing bradycardia, 17
 - causing tachypnea, 20
 - dosage rate, 330
 - mild colic, 46
 - postoperative pain, 250–251
 - postoperative patient care, 239
- diaphragmatic hernia
 - laparoscopy, 177, 188
 - postpartum mares, 285
 - radiography, 157
 - sonographic evaluation, 136, 138
- diarrhea *see also* enterocolitis; colitis
 - cyathostomins, 320
 - enteral fluid therapy, 68
 - enterocolitis, 279
 - in postoperative colics, 231
 - IV fluid therapy
 - acidosis, 111
 - electrolyte abnormalities, 110
 - maintenance fluid rate, 110
 - postoperative
 - prevalence, 248
 - prevention, 248
 - treatment, 248–9
 - small colon impaction, 221–2
 - ventral midline celiotomy, 248–9
- dimethyl sulfoxide, 53, 330
- dioctyl sodium sulfosuccinate (DSS), 330
 - laxative, 66
 - mild colic, 47
- early referral *see* referral
- electrolytes
 - abnormality
 - hypernatremia, 68, 110
 - hypocalcemia, 68, 82, 111
 - hypochloremia, 68, 82
 - hypokalemia, 82
 - hypomagnesemia, 68, 82
 - hyponatremia, 65, 109, 110
 - parenteral nutrition, 310
 - enteral fluid therapy, 68
 - imbalance, 81–2
 - IV fluid therapy, 110–111
- endotoxemia
 - electrolyte imbalances, 82, 111
 - fluid distribution, 100
 - fluid therapy, 111–14
 - hetastarch, 113
 - hypertonic saline solution, 112
 - hypertriglyceridemia, 288
 - injected oral mucous membranes, 17–18
 - lactate concentration, 81
 - leukopenia, 80
 - lidocaine (IV), 57

- endotoxemia (*cont'd*)
 - NSAID, 52
 - platelets, 83–4
 - pregnant mares, 283
 - prevention, 246–7
 - pyrexia, 19
 - secondary hemostatic system, 84
 - shock, 73, 101
 - treatment, 247–8
- enrofloxacin, 331
 - colic surgery, 175
 - dose rate, 331
 - postoperative patient care, 240
- enteral fluid therapy
 - complications
 - diarrhea, 68
 - electrolyte disturbances, 68
 - mild colic, 67–8
 - nasogastric tube-associated problems, 68–9
 - contraindications, 64
 - enteral fluids composition
 - balanced electrolyte solution, 65
 - dioctyl sodium sulfosuccinate (DSS), 66
 - laxatives or cathartics, 66
 - magnesium sulfate, 66
 - mineral oil, 66
 - oral rehydration solutions, 65
 - tap water, 64–5
 - fluids and electrolytes absorption, 62–3
 - indications, 63–4
 - technique
 - continuous infusions, 66–7
 - intermittent boluses, 66
- enterocentesis
 - abdominocentesis, 89
 - peritoneal fluid analysis, 90
- enterocolitis *see also* diarrhea; colitis
 - decreased AT activity, 85
 - electrolyte abnormalities, 82
 - leukopenia, 80
 - necrotizing enterocolitis
 - sonographic evaluation, 120
 - neonatal foals, 4, 279–80
 - postoperative patient care, 231
 - triglyceride abnormalities, 83
- enterolithiasis, 218–9
 - diet, 7, 296
 - palpation per rectum, 27, 28
 - radiography, 155–6
 - recurrent colic, 9, 73
 - risk factors, 218
 - signs, 219
 - sonography, 141
 - treatment of, 219
 - water source, 8, 297
- epiploic foramen entrapment, 211
- equine grass sickness, 209
- erythrogram, 79
- esophageal trauma, NGT, 44
- exploratory celiotomy *see* colic surgery

- fast localized abdominal sonography of horses (FLASH), 118
- fecal egg count reduction tests (FECRT), 317
- fecal egg counts (FECs), 317
- FECRT *see* fecal egg count reduction tests (FECRT)
- FECs *see* fecal egg counts (FECs)
- fenbendazole, 209, 319, 321–3, 333
- fever
 - biosecurity, 270
 - geriatric and mature horses, 287
 - postoperative complication, 244–6
 - postoperative patient care, 232
 - rectal temperature, 19
- firocoxib, 331 *see also* non-steroidal anti-inflammatory drugs (NSAID)
 - clinical use, 53
 - ischemia-reperfusion injury, 246
 - mechanism of action, 52
 - mild colic, 46
 - postoperative pain, 250
 - postoperative patient care, 239
- FLASH *see* fast localized abdominal sonography of horses (FLASH)
- fluid distribution, 99–101
 - colloid oncotic pressure, 100–101
 - dehydration, 101
 - endotoxemia and hypoproteinemia, 100
 - hypovolemia, 101
 - Landis-Pappenheimer equation, 100
 - Starling-Landis equation, 100
- fluid therapy
 - enteral (*see* enteral fluid therapy)
 - IV (*see* intravenous (IV) fluid therapy)
 - mild colic, 47
- flunixin meglumine, 6, 36, 57, 72, 83, 331, 333 *see also* non-steroidal anti-inflammatory drugs
 - adhesion prevention, 255, 256
 - adverse effects and contraindications, 53
 - clinical use, 52

- endotoxemia (treatment), 247, 249
- indications
 - for referral, 72
 - for surgery, 165–6
- mechanism of action, 52
- mild colic, 46–7
- postoperative pain, 250
- postoperative patient care, 238–9
- preparation for surgery (analgesia), 174
- foals, 279–81
 - adhesions, 255–6
 - aminoglycoside plasma concentration, 339
 - abdominocentesis
 - omental herniation, 89
 - biochemistry, 338
 - causes of colic, 279–81
 - ascarid impaction, 208–9
 - cecal impaction, 214
 - gastric ulceration, 205
 - jejunal volvulus, 212
 - meconium retention, 222–3
 - colloid oncotic pressure (COP), 100
 - colic, 278–9
 - management, 283–4
 - neonates, 279–80
 - postoperative complication, 283
 - young foals and weanlings, 280–281
 - colic surgery
 - incision, 177
 - needle decompression, 189
 - preparation, 175
 - drug dosages, 329–35
 - energy and protein requirements for, 302
 - fluid therapy, 109–111
 - gamma glutamyl transferase (GGT), 82
 - hematology, 337
 - IV catheterization, 103–9
 - mean arterial blood pressure, 102
 - nasogastric tube, 39, 40
 - nutrition
 - energy requirements, 302
 - enteral nutrition, 304–5
 - general, 302
 - parenteral nutrition, 308–10
 - protein requirements, 303
 - Parascaris equorum*, 321–2
 - peritoneal fluid
 - nucleated cell count, 93
 - physical examination
 - heart rate, 16
 - rectal temperature, 19
 - prognosis for racing, 295
 - radiography, 155, 157–9
 - sonography, 46, 117, 119–20
 - abscess, 138–40
 - jejunojejunal intussusception, 127
 - meconium retention or impaction, 125
 - stomach, 132
- gamma glutamyltransferase (GGT), 82–3
- gas colic, 4, 9, 45, 49 *see also* mild colic; spasmodic colic
- gastric disease
 - gastric impaction, 205–6
 - gastric rupture, 206–7
 - gastric ulceration, 205
- gastrointestinal parasitology
 - Anoplocephala perfoliata*, 318–19
 - anthelmintics, 319, 323
 - cyathostomes, 320–321
 - FECs and FECRT, 317
 - Parascaris equorum*, 321–22
 - parasite control programs, 317–18
 - parasites, 316–17
 - Strongylus vulgaris*, 322–3
- gastrosplenic ligament entrapment, 211–12
- gentamicin, 162, 240, 240, 332
- geriatric horses, 286–7
- heart rate
 - assessment, 16–17
 - hyoscine-N-butylbromide, 59
 - packed cell volume, 79
 - postoperatively, 232
 - shock, 73
- hemorrhage
 - effusions, 96, 97
 - epiploic foramen entrapment, 211
 - hemangiosarcoma, 142
 - intra-abdominal, 134
 - intraluminal uterine, 136
 - IV fluid therapy, 109
 - mesenteric trauma, 222
 - nasogastric intubation, 43–4
 - pale mucous membranes, 18
 - peritoneal fluid, 90, 97
 - periparturient, 135
 - postoperatively, 233
 - postpartum mares, 34, 170, 284
 - pregnant mares, 284

- hemorrhage (*cont'd*)
 - shock, 101
 - trocharization, 163
- hepatobiliary assessment, 82–3
- hetastarch, 108, 113–14, 332
 - postoperative patient care, 236–7
 - preparation for colic surgery, 174
- Holliday-Segar formula, 110
- HSS *see* hypertonic saline solution (HSS)
- hyoscine-N-butylbromide *see also* Buscopan®
 - adverse effects and contraindications, 59
 - clinical use, 58–9
 - mechanism of action, 58
- hyperglycemia, 83
 - parenteral nutrition, 309–10
 - postoperative patient care, 234
- hyperlipemia
 - parenteral nutrition, 310
 - ponies, miniature horses, and donkeys, 288–9
- hypermagnesemia, enteral fluid therapy, 68
- hypernatremia
 - electrolyte disturbances, 68
 - IV fluid therapy, 110
- hypertonic saline solution (HSS), 111–12, 235, 248, 332
- hypertriglyceridemia
 - parenteral nutrition, 309–10
 - ponies, miniature horses, and donkeys, 288–9
 - postoperative patient care, 334
 - triglycerides, 83
- hypocalcemia, 82
 - enteral fluid therapy, 68
 - IV fluid therapy, 111
- hypochloremia, 68, 82
- hypoglycemia, 83
- hypokalemia, 82
- hypomagnesemia, 68, 82
- hyponatremia
 - colitis, 216
 - enteral fluids, 65, 68
 - IV fluid therapy, 109, 110
- ileal impaction, 208
 - Anaplocephala perfoliata*, 9, 318
 - geographical regions, 8
 - indications for surgery, 167
 - palpation per rectum, 27, 28
 - prevention, 296
 - sonography, 130
 - surgery, 199
 - tapeworms, 9, 318
- ileal muscular hypertrophy *see* idiopathic small intestinal muscular hypertrophy
- idiopathic small intestinal muscular hypertrophy, 129, 130, 133, 151, 208
- ileocecal intussusceptions, 213–14
 - Anoplocephala perfoliata*, 9, 10, 318
 - palpation per rectum, 27
 - signalment, 4
 - sonography, 127, 129
 - surgery, 199
 - tapeworms, 9, 10, 318
- ileocecostomy, 194–5
- impaction colic *see* ileal impaction; cecal impaction; large colon impaction; small colon impaction
- incisional complications, 256–8
- inguinal hernia, 28, 29, 199
- intestinal borborygmi, 19
- intestinal obstruction
 - complete/partial intestinal, 165–6
 - early referral, 73–4
 - hypochloremia, 82
 - small intestine, 131
- intestinal resection and anastomosis, 192
 - ileocecostomy, 194–5
 - jejunocecostomy, 193–4
 - jejunoileostomy, 193
 - jejunojejunostomy, 193
 - large colon resection
 - end-to-end technique, 195–7
 - side-to-side anastomosis, 197
 - small colon resection, 197–202
- intra-abdominal or intra-peritoneal adhesions *see* adhesions
- intravenous (IV) catheterization
 - complications, 107, 109
 - indications, 103
 - preparation
 - catheters, 103–4
 - materials required for, 103
 - skin preparation, 104
 - procedure
 - IV catheter care, 106–7
 - over-the-needle technique, 105–6
 - over-the-wire (Seldinger) technique, 106–7
- intravenous (IV) fluid therapy
 - colloid replacement
 - hydroxyethyl starch, 113–14
 - plasma, 112
 - hypertonic saline solution (HSS), 111–12

- oxygen delivery
 - central venous pressure, 102
 - hypovolemia, 101
 - hypoxia, 102
 - mean arterial blood pressure, 102–3
 - plasma creatinine concentration, 102
 - shock, 101–2
 - SvO₂ and venous oxygen tension, 102
 - urine output, 102
- polyionic isotonic fluids
 - electrolyte replacement, 110–111
 - maintenance fluid rate, 110
 - replacement/resuscitation, 109–10
- intravenous (IV) fluids, 108
- intravenous (IV) lidocaine, 57–8 *see also*
 - analgesia
 - clinical use, 57–8
 - complications, 58
 - dose rate, 332
 - ischemia-reperfusion injury, 246
 - mechanism of action, 57
 - postoperative ileus, 251–3
 - postoperative pain, 250
 - postoperative patient care, 239
- ischemia-reperfusion injury, 246
 - adhesions, 255
 - dimethylsulfoxide (DMSO), 330
 - early referral, 71, 72
 - flunixin meglumine and, 53
 - hypocalcemia, 82, 111
 - hypomagnesemia, 82, 111
 - ileus, 251
 - intravenous (IV) lidocaine and, 57–8, 239
 - lactate, 79, 81, 92
 - leukopenia, 80, 245
 - magnesium sulfate, 111
 - postoperative complication, 246
- intussusception *see* ileocecal intussusception;
jejunojejunal intussusception; cecocolic
intussusception
- ivermectin-based anthelmintics, 9, 209, 296, 297,
319, 320–322, 332
- jejunal intussusception, 213
- jejunal volvulus, 4, 14, 27, 180, 194, 198, 212, 246,
279–80, 322
- jejunocecostomy, 193–4
- jejunoileostomy, 193
- jejunojejunal intussusception, 4, 27, 127, 128, 181,
209, 210, 213, 279–80, 322
- jejunojejunoscopy, 193
- jugular refill time, 18
- ketamine, 239, 250, 332, 334
- kidney
 - palpation per rectum, 25, 26
 - nephrosplenic ligament entrapment, 29, 31,
35, 219
 - non-steroidal anti-inflammatory drugs,
53, 239
 - sonographic evaluation, 118–20
 - nephrosplenic ligament entrapment, 121–2
 - neoplasia, 141
 - urolithiasis, 143–5
 - surgical anatomy, 183, 186, 188
- laminitis, 5, 20–21, 169, 208, 216, 232, 233, 248, 255,
292, 295
- laparoscopy, 37, 177, 187–8, 199–201, 212, 255, 256,
258, 285, 294
- large colon disease
 - abdominal palpation per rectum, 28–9, 32–3
 - inflammatory disease, 215–6
 - simple obstruction
 - ascending colon impaction, 216
 - enterolithiasis, 218–19
 - nephrosplenic ligament entrapment, 219–20
 - right dorsal displacement, 220
 - sand colopathy, 217–18
 - sonographic evaluation
 - large colon volvulus, 122–3
 - left dorsal displacement, 121–2
 - right dorsal displacement, 120–121
 - strangulating obstruction, 220–221
- large colon volvulus (LCV), 220–221, 324
 - clinical signs, 6
 - CVP measurement, 102
 - history, 9
 - palpation per rectum, 32
 - plasma lactate concentration, 170–171
 - postpartum mare, 4, 170, 284, 285
 - prognostic indices, 170
 - referral, 72
 - sonographic evaluation, 120–123
 - surgery, 195, 221
- Lawsonia intracellularis* infection, 281
- laxatives, 47, 66
- LCV *see* large colon volvulus (LCV)
- left dorsal displacement *see* nephrosplenic
ligament entrapment leukogram, 79–80, 337

- lidocaine *see also* intravenous lidocaine
 enema
 for palpation per rectum, 23–4
 for rectal tear, 36
 intravenous catheterization, 103, 106
 nasogastric intubation, 42
 trocharization, 161
- lipopolysaccharide *see* endotoxin
- Magnesium, 338
 hypermagnesemia, 68
 hypomagnesemia, 68, 82
 IV fluid therapy, 111
- magnesium sulfate, 333
 enteral fluid therapy, 66
 ischemia-reperfusion injury, 111
 mild colic, 47
- mean arterial blood pressure (MAP), 102–103
- Meckel's diverticulum, 212–13
- meconium-associated colic, 222–3, 279–80
 radiographic evaluation, 158
 signalment, 4
 sonographic evaluation, 120, 125
- meloxicam, 53 *see also* non-steroidal anti-inflammatory drugs
 ischemia-reperfusion injury, 246
 mild colic, 46
 postoperative patient care, 239
- mesenteric rent, 199, 222
- mesodiverticular band, 213
- metabolic indicators, 83
- mild colic
 clinical indications, 45–6
 enteral fluid therapy, 67–8
 management
 analgesia, 46–7
 fluid therapy, 47
 frequent monitoring, 48
 laxatives, 47
 nutrition, 48
 spasmolytic drugs, 47
 practical perspective, 45
- miniature horses, 288–9
- N*-butylscopolammonium bromide *see* Buscopan®
- nasogastric intubation *see also* nasogastric tube (NGT)
 complications
 hemorrhage, 43–4
 pharyngeal and esophageal trauma, 44
 tracheal intubation, 44
 early referral, 75
 gastric impaction, 43
 indications, 38
 preparation
 buckets and water, 39
 head-neck ventroflexion, 39
 materials required, 38, 39
 medications, 39
 nasogastric tube, 39, 40
 short-acting sedative, 39
 stomach pump, 39
 procedure
 nasogastric tube passage, 40–42
 reflux assessment, 42–3
 sedation, 39–40
 proximal enteritis, 43
- nasogastric tube (NGT) *see also* nasogastric intubation
 passage of
 esophagus, 41
 nasal passage, 41
 pharynx, 41
 stomach, 41–2
 trachea, 41–2
 permanent marks, 39–40
 problems with, 44
 types of, 39–40
- necrotizing enterocolitis, neonatal foals, 279–80
- needle technique, abdominocentesis, 88–9
- neonatal foals *see also* foals
 causes of colic, 279
 colloid oncotic pressure (COP), 100
 fluid therapy, 110–111
 gamma glutamyltransferase (GGT), 82
 hematology in, 337
 management, 281
 meconium retention, 222
 mixed venous oxygen, 102
 plasma antimicrobial drug concentrations, 339
 plasma biochemistry for, 338
 sonographic evaluation, 117, 119–20, 132
- nephrosplenic ligament entrapment (NSLE), 183, 219–20
 abdominal palpation per rectum, 29, 32
 diagnosis, 219
 sonographic diagnosis, 121–2
 surgical correction, 200
 prevention, 294

- nonsteroidal anti-inflammatory drugs
(NSAIDs)
adverse effects and contraindications, 53–4
clinical use, 52–3
mechanism of action, 52
mild colic, 46
postoperative patient care, 238–9
for young foals and weanlings, 281
- nostril flare, 19–20
- NSAIDs *see* nonsteroidal anti-inflammatory drugs
(NSAIDs)
- NSLE *see* nephrosplenic ligament entrapment
(NSLE)
- omental herniation, 89–90
- opioids
adverse effects and contraindications, 56–7
analgesic therapy, 46
antagonists, 56
clinical use, 56
mechanism of action, 55
- oral mucous membranes
dry, 18
hemorrhagic shock, 18
injected membranes, 17–18
normal, 17
tacky, 18
- over-the-needle technique, 105–6
over-the-wire (Seldinger) technique, 106–7
- Parascaris equorum*, 9, 321–2 *see also* ascariasis
- partial typhlectomy, 191–2
- patient's history
clinical presentation, 323–7
clinical significance, 1–3
detailed history
appetite, 6–7
defecation and urination, 7
exercise regimen, 8
feeding regimen, 7
gastrointestinal parasite control, 8–9
housing location, 8
pasture access and housing, 8
recent transportation, 8
vaccination history, 9
water consumption, 7
water source, 7–8
- initial history
clinical signs, 6
colic signs and duration, 6
dull mentation and inappetence, 5–6
mare's reproductive status, 6
recumbency, 5
- medical history
crib biting/ windsucking, 10
current/recent medication, 9–10
previous colic and colic surgery, 9
recent medical problems, 10
- signalment, 4
- pelvic flexure (PF) impactions, 217
enteral fluid therapy, 63–8
indications for surgery, 167
medical history, 6, 7, 9–10
palpation per rectum, 32–3
pelvic flexure enterotomy, 190
physical examination, 14
signalment, 4
sonographic evaluation, 123
- pelvic flexure enterotomy, 190
- pentastarch, 113, 114
- peritoneal cavity
abdominal palpation per rectum, 35–6
hematoma, 134, 135
- peritoneal fluid analysis
application of, 97
biochemical evaluation
D-dimer concentrations, 92
lactate and glucose concentration, 92
protein concentration, 91–2
- cells and cell counts
erythrocytes, 93
nucleated cells, 93–5
- effusion
exudates, 97
hemorrhagic, 97
transudate, 95–7
- gross characteristics
color and clarity, 90–91
volume, 90
- indications, 87–8
nonrepresentative sampling, 90
- personal protective equipment (PPE), 266
- pharyngeal trauma, nasogastric intubation, 44
- physical examination
admission physical examination sheet, 13
cardiovascular status assessment
dehydration, hypovolemia, and shock,
15–16
extremity temperature, 18
heart rate, 16–17

- physical examination (*cont'd*)
- jugular refill time, 18
 - oral mucous membranes, 17–18
 - pulse quality, 19
 - checklist, 14
 - clinical significance, 12–14
 - digital pulses, 20–21
 - gastrointestinal tract evaluation, 19
 - initial patient observation
 - abdominal distention, 14
 - abrasions, 14
 - body condition score, 14–15
 - mentation/attitude, 14
 - pain assessment, 14–15
 - sweating, 14
 - rectal temperature, 19
 - respiratory system, 19–20
- piperazine, 209, 319, 336
- plasma, 112
- pneumoperitoneum, 36
- polyurethane catheters, 106, 107
- ponies, 287–8
- postoperative complications
 - diarrhea, 248–9
 - endotoxemia and SIRS, 246–8
 - fever, 244–6
 - incisional complications, 256–8
 - intraperitoneal adhesion formation, neonates, 281
 - intraperitoneal adhesions, 255–6
 - ischemia-reperfusion injury, 246
 - laminitis, 20–21, 295
 - postoperative ileus, 251–3
 - recurrent colic, 249–51
 - septic peritonitis, 253–4
 - small intestinal volvulus, 212
- postoperative ileus, 251–3
- postoperative intraperitoneal adhesions, 255–6
- postoperative patient care
 - monitoring
 - arterial blood pressure, 234–5
 - central venous pressure, 234
 - laboratory data, 233–4
 - patient assessment, 231–2
 - physical examination, 232–3
- treatment
 - anti-inflammatory and analgesic drugs, 238–40
 - antimicrobial drugs, 239–41
 - colloids, 236
 - crystalloids, 235–6
 - fluid therapy, 235, 237–8
 - hydroxyethyl starch, 236–7
 - plasma, 236
 - postoperative feeding and nutrition, 241
- postpartum mares, 284–5
- potassium supplementation, 110
- PPE *see* personal protective equipment (PPE)
- praziquantel, 9, 208, 214, 319, 320, 336
- pregnant mares, 282–4
- prevention of colic, 295–8
- proliferative enteropathy *see* *Lawsonia intracellularis* infection
- prostanoids, 52
- proximal enteritis, 207–208
 - gamma glutamyltransferase, 82–3
 - indications for surgery, 167
 - mild colic, 49
 - nasogastric reflux, 43, 73
 - sonographic evaluation, 129–30
- psyllium mucilloid, 47, 297, 334
- pulse quality, 19
- PvO₂ *see* venous oxygen tension (PvO₂)
- Radiography *see* abdominal radiography
- RDD *see* right dorsal displacement (RDD)
- rectal examination *see* abdominal palpation per rectum
- rectal tear, 34
 - emergency first aid, 36
 - management, 37
- recurrent colic, 9, 14, 46, 72–4, 143, 165, 170, 223, 249–51, 294–8
- referral
 - client
 - financial considerations, 76
 - referral hospital, 75–6
 - clinical significance, 71–2
 - indications
 - abdominal pain, 72–3
 - abdominal palpation per rectum, 74
 - intestinal obstruction, 73–4
 - nasogastric reflux, 73
 - shock, 73
 - tachycardia/ increasing heart rate, 73
 - patient preparation
 - analgesia, 74–5
 - fluid therapy, 75
 - nasogastric intubation, 75
 - transportation, 75
 - trocharization, 75
- Rhodococcus equi* intra-abdominal abscess, 140
- right dorsal colitis, 216–17

- medical history, 9
- medical *vs.* surgical treatment, 169
- non-steroidal anti-inflammatory drugs (NSAIDs), 53
- sonographic evaluation, 125–6
- surgery, 188
- right dorsal colon (RDC) impactions, 33 *see also*
 - large colon impaction
- right dorsal displacement (RDD), 220
 - colonic impaction and, 168
 - gamma glutamyltransferase, 82
 - medical history, 9
 - palpation per rectum, 28–9, 32, 72
 - postpartum mares, 284
 - sonographic evaluation, 120–121
 - surgical correction, 200
- SDH *see* sorbitol dehydrogenase (SDH)
- scrotal hernia, 137
- septic peritonitis
 - cecal perforation, 215
 - postoperative patient care, 232
 - postoperative complication, 253–4
- SIRS *see* endotoxin
- SISO *see* small intestinal strangulating obstruction (SISO)
- small (descending) colon
 - abdominal palpation per rectum, 33
 - small (descending) colon impaction, 221–2
 - meconium retention, 222–3
 - mesenteric rent, 222
 - neoplasia, 223
- small colon enterotomy, 191
- small intestinal enterotomy, 191
- small intestinal strangulating obstruction (SISO), 209–14
 - clinical pathology, 80, 82–3
 - neonates, 279
 - postoperative feeding, 306
 - referral, 71
- sonography *see* abdominal sonographic evaluation
- sorbitol dehydrogenase (SDH), 82, 338
- spasmodic colic, 205, 306, 311, 316, 318 *see also* gas colic; mild colic
- spasmolytic drugs, 47 *see also* Buscopan®
- spleen
 - abdominal palpation per rectum, 25, 26, 35
 - nephrosplenic ligament entrapment, 29, 31
 - gastrosplenic ligament entrapment, 211–12
 - neoplasia, 223
 - nephrosplenic ligament entrapment, 219
- sonographic evaluation, 119, 120
 - abscess, 138, 139
 - adhesions, 130
 - gastric distention, 131, 132
 - hematoma, 140
 - hemorrhage, 134
 - nephrosplenic ligament entrapment, 121–2
 - neoplasia, 141
 - surgical anatomy, 183, 186, 188
- strangulating *vs.* nonstrangulating obstruction, 128–30, 166–7
- Strongylus vulgaris*, 9, 322–3
- surgical treatment *see also* colic surgery
 - euthanasia without surgery, 170–171
 - gastrointestinal tract perforation, 171
 - prognostic indices, 170–171
- indications for
 - cecal impaction, 168
 - colitis, 169
 - intestinal obstruction, 165–6
 - large colon impaction, 167–8
 - medical management response, 165
 - peritoneal fluid analysis, 166
 - peritonitis, 169–70
 - persistent or severe abdominal pain, 165
 - postpartum hemorrhage, 170
 - small colon impaction, 168–9
 - small intestinal lesions, 166–7
 - sonographic examination, 166
- SvO₂ *see* venous oxygen saturation (SvO₂)
- systemic inflammatory response syndrome *see* endotoxemia
- tachycardia, 16–8, 73, 167, 232, 323–5
 - heart rate assessment, 16, 17, 232
- tachypnea, 20, 55, 232
 - respiratory rate assessment, 20
- tapeworms *see* *Anoplocephala perfoliata*
- teat cannula technique for abdominocentesis, 89
- transudates, 95–7
- trocharization
 - complications, 162–3
 - indications, 160–161
 - preparation, 161
 - prior to referral, 75
 - procedure
 - paralumbar approach, 161–2
 - transrectal approach, 162
- typhlotomy, 190–191

- ultrasonography, 46, 121–2 *see also* abdominal sonographic evaluation
- umbilical hernia, 138
- urogenital tract
 - abdominal palpation per rectum, 34–5
 - transrectal examination, 120
- uterine hemorrhage, 136–7, 284–5
- uterine torsion, 34–5, 282

- venous oxygen saturation (SvO_2), 101, 102
- venous oxygen tension (PvO_2), 101, 102
- ventral midline celiotomy
 - approach, 176–8
 - closure, 183–5

- xylazine, 54–5, 335
 - bradycardia, 17
 - colic surgery preparation, 175
 - epidural, 36
 - indications
 - for referral, 72
 - for surgery, 165
 - mild colic, 46–7
 - motility, 63, 117,
 - nasogastric intubation, 39
 - palpation per rectum, 24
 - postoperative pain, 250
 - postoperative patient, 239
 - rectal tear, 36
 - tachypnea with fever, 20